



Winter Braids Lecture Notes

Participants & Programme

Vol. 5 (2018), p. i-vi.

<http://wbln.centre-mersenne.org/item?id=WBLN_2018_5_r1_0>

cedram

Texte mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.centre-mersenne.org/>

Participants & Programme

This volume of *Winter Braids Lecture Notes* contains the lecture notes for the four mini-courses given at Winter Braids VIII, which took place at the CIRM in Marseille from February 5th to 9th, 2018. This eight edition of the school was part of the thematic month *Low Dimensional Topology and Related Topics* at CIRM.

We thank Benjamin Audoux for his great work as local organizer for this edition of the Winter Braids school.

Participants

Amiot	Claire	(Université Grenoble Alpes)
Audoux	Benjamin	(Université Aix-Marseille)
Baader	Sebastian	(Universität Bern)
Bagherifard	Narges	(Sharif University Iran)
Bagherifard	Neda	(Amirkabir University Iran)
Bagnoud	Pierre Aldéric	(Université de Genève)
Bahayou	Amine	(Université Kasdi Merbah)
Bakshi	Rhea Palak	(George Washington University)
Bamba	Klanan Bertrand	(Université de Cocody)
Baranowski	Adam	(University of Cambridge)
Belletti	Giulio	(Scuola Normale Superiore di Pisa)
Bellingeri	Paolo	(Université de Caen Normandie)
Ben Aribi	Fathi	(Université de Genève)
Berat	Clément	(Université Grenoble Alpes)
Bode	Benjamin	(University of Bristol)
Boileau	Michel	(Aix-Marseille Université)
Bonnier	virginie	(Université de Genève)
Brandenburgsky	Michael	(Ben Gurion University)
Brendle	Tara	(University of Glasgow)
Casejuane	Adrien	(Université Grenoble Alpes)
Caudell	Jacob	(Boston College)
Celoria	Daniele	(University of Oxford)
Cisneros	Bruno	(Conacyt Imaca Oaxaca)
Cumplido Cabello	Maria	(Université de Rennes 1)
Damiani	Celeste	(Osaka City University)
Dey	Soumya	(IISER Mohali India)
Diarrassouba	Siriky	(Université de Cocody)
Duran ?Batalla	Juan Luis	(Universidad Autonoma Barcelona)
Dvirnas	Albertas	(Lund University)
Estier	Solenn	(Université de Genève)
Faitg	Matthieu	(Université de Montpellier)
Feller	Peter	(ETH Zürich)
Florens	Vincent	(Université de Pau)

Participants & Programme

Földvári	Viktória	(Eötvös Loránd University)
Fossati	Edoardo	(SNS PISA)
Gadbled	Agnès	(Uppsala Universitet)
Gandolfi	Guillaume	(Université Caen-Normandie)
Gay	David	(University of Georgia)
Gonzalez	Pagotto	(Université Grenoble Alpes)
Gultepe	Funda	(University of Luxembourg)
Haissinsky	Peter	(Aix-Marseille Université)
Hok	Jean-Marc	(Université Toulouse III)
Ibrahim	Abdoulrahim	(Université de Montpellier)
Kegel	Marc	(University of Cologne)
Kjuchukova	Alexandra	(University of Wisconsin Madison)
Kolay	Sudipta	(Georgia Institute of Technology)
Kosanovic	Danica	(Max Planck Institute, Bonn)
Kulkarni	Ajinkya	(Université de Bourgogne)
Lange	Christian	(University of Cologne)
Lee	Sang-jin	(Konkuk University)
Lee	Eon-Kyung	(Sejong University)
Leturcq	David	(Université Grenoble Alpes)
Liechti	Livio	(Institut de Math. de Jussieu)
Limouzineau	Maÿlis	(University of Cologne)
Lopez	Daniel	(Université Paris Diderot)
Martel	Jules	(Université Toulouse III)
Masbaum	Gregor	(Université Pierre et Marie Curie)
McLeay	Alan	(University of Glasgow)
Meier	Jeffrey	(University of Georgia)
Meilhan	Jean-Baptiste	(Université Grenoble Alpes)
Meumertzheim	Johanna	(University of Regensburg)
Mihajlovic	Stefan	(Central European University)
Misev	Filip	(Université Aix Marseille)
Moussard	Delphine	(RIMS, Kyoto University)
Owad	Nicholas	(Okinawa Institute of Sciences and Technology)
Palesi	Frédéric	(Université Aix Marseille)
Palmer-Anghel	Cristina	(Université Paris Diderot)
Palmer-Anghel	Martin	(Universität Bonn)
Paoluzzi	Luisa	(Université Aix Marseille)
Pichon	Anne	(Université Aix Marseille)
Pinochet Lobos	Antoine	(Université Aix Marseille)
Rezaeenayeh	Khadijeh	(Tehran polytechnic)
Robert	Louis-Hadrien	(Université de Genève)
Sane	Abdoul Karim	(ENS Lyon)
Silvero	Marithania	(Inst. of Math. Polish Academy)
Singh	Oliver	(University of Durham)
Thiel	Anne-Laure	(Stuttgart Universität)
Thomas	Alexander	(Université de Strasbourg)
Tovstopyat-Nelip	Lev	(Boston College)
Ueki	Jun	(The University of Tokyo)
Vera	Anderson	(Université de Strasbourg)
Vershinin	Vladimir	(Université de Montpellier)
Viana Bedoya	Natalia	(Universidade Federal de Sao Carlos)
Wagner	Emmanuel	(Université de Bourgogne)
Walsh	Genevieve	(Tufts University)
Witzel	Stefan	(Bielefeld University)
Zuddas	Daniele	(Universität Bayreuth)

Abstracts of Courses

Claire Amiot (Univ. Grenoble Alpes)

Cluster algebras from surfaces and categorification

In this course I will first introduce cluster algebras associated with a triangulated surface. I will then focus on representation of quivers, and show the strong link between cluster combinatorics and representation theory. The aim will be to explain additive categorification of cluster algebras in this context. All the notions will be illustrated by examples.

Michel Boileau (Univ. Aix-Marseille)

3-manifolds groups and related topics

Recent years have seen spectacular progress in the understanding of the algebraic properties of the fundamental groups of 3-manifolds. In these lectures we will present some of these results. We will start with some basic properties of 3-manifold groups related to topology. Then we will discuss the relationship with geometric properties of 3-manifolds, with an emphasis on surface subgroups. The last lecture will focus on virtual properties and profinite properties of 3-manifold groups.

Tara Brendle (Univ. Glasgow)

Congruence braid groups

This course will focus on braid groups, as viewed through the lens of a symplectic representation. We will discuss various contexts in which this representation arises, including the classical symplectic representation of mapping class groups and the Burau representation. The kernel of this representation, often referred to as the braid Torelli group, also admits a number of different definitions including an algebro-geometric description. Along the way, we will also describe generalizations of the braid Torelli group, such as the Torelli group for an arbitrary surface and braid congruence groups, and we will give a sample application to the Brunnian braid group.

David Gay (Univ. Georgia)

From Heegaard splittings to trisections; porting 3-dimensional ideas to dimension four

Trisections are to 4-manifolds as Heegaard splittings are to 3-manifolds. I will develop the foundations of the theory (why do trisections exist and how unique are they?) and then survey a range of newer extensions of the theory (submanifolds, boundaries, gluing). Throughout I will be emphasizing what we don't know and advertising open problems; with luck, maybe some will be solved by participants and we'll all know more at the end of the week!

Abstracts of Short Talks

Cristina Anghel-Palmer (Univ. Paris 7)

A homological model for the coloured Jones polynomials

In 1991, Reshetikhin and Turaev defined a method that starts with a quantum group and leads to link invariants. This construction is purely algebraic and combinatorial. The coloured Jones polynomials $J_N(L, q)$ are a family of quantum invariants constructed from the representation theory of $U_q(\mathfrak{sl}(2))$. We will describe a geometrical interpretation for the coloured Jones polynomials.

R. Lawrence defined a sequence of representations of the braid groups, using the homology of a certain covering of a configuration space. In 2012, Kohno proved a deep connection between the representations of the braid group on the highest weight spaces of $U_q(\mathfrak{sl}(2))$ -modules and the Lawrence representations. Using this, we give a homological model for $J_N(L, q)$. We prove that the coloured Jones polynomials can be described as a graded intersection pairing between two homology classes on a covering of the configuration space of the punctured disc.

Maria Cumplido Cabello (Univ. Rennes 1 / Univ. Sevilla)

Advances in the understanding of parabolic subgroups of Artin-Tits groups

Artin-Tits groups are a natural generalisation of braid groups from the algebraic point of view: In the same way that the braid group can be obtained from the presentation of the symmetric group with transpositions as generators by dropping the order relations for the generators, other Coxeter groups give rise to more general Artin-Tits groups. If the underlying Coxeter group is finite, the resulting Artin-Tits group is said to be of spherical type. Artin-Tits groups of spherical type share many properties with braid groups.

The properties of the complex of curves, and the way in which the braid group acts on it, allow to use geometric arguments to prove results in braid groups. As one cannot replicate these geometrical techniques in other Artin-Tits groups, they must be replaced by algebraic arguments if one tries to extend properties of braid groups to all Artin-Tits groups of spherical type. That is why we are interested in parabolic subgroups of Artin-Tits groups, which are defined as conjugates of a subgroups generated by a subset of the standard generators. They are the analogue of isotopy classes of simple closed curves in the puncture disk, which are the building blocks that form the complex of curves. Then, it is logical to believe that improving our understanding about parabolic subgroups will allow us to prove similar results for Artin-Tits groups of spherical type in general.

In this talk two new results about parabolic subgroups will be presented, namely that the intersection of parabolic subgroups is a parabolic subgroup and that the set of parabolic subgroups is a lattice.

Joint work with Volker Gebhardt, Juan González-Meneses and Bert Wiest.

Peter Feller (ETH Zurich)

Braids with as many full twists as strands realize the braid index

We explain how the fractional Dehn twist coefficient of braids—a rational quantity studied in mapping class group theory—can be recast in terms of a certain linear combination of (concordance) invariants of knots. This interpretation allows to provide lower bounds on the braid index of closures of braids with “many twists”.

Based on joint work with Diana Hubbard.

Participants & Programme

Marc Kegel (Univ. Köln)

The knot complement problem for Legendrian and transverse knots

The famous knot complement theorem by Gordon and Luecke states that two knots in the 3-sphere are equivalent if and only if their complements are homeomorphic. In this talk I want to discuss the same question for Legendrian and transverse knots and links in contact 3-manifolds. The main results are that Legendrian as well as transverse knots in the tight contact 3-sphere are equivalent if and only if their exteriors are contactomorphic.

Alexandra Kjuchukova (Univ. Wisconsin Madison)

Trisections of singular branched covers

Let Y be a closed oriented four-manifold which is an irregular dihedral cover of the four-sphere, branched along an oriented surface B embedded in S^4 with one cone singularity. I will describe a method to produce a trisection of Y from a singular triplane diagram of the pair (S^4, B) . I will give a construction of an infinite family of three-fold dihedral covers $f_i : Y_i \rightarrow S^4$, branched along singularly embedded two-spheres whose singularities are cones on pairwise distinct knot types K_i . I will use trisections to prove that, for infinitely many among these knots K_i , the manifold Y_i is diffeomorphic to CP^2 . Lastly, I will discuss a ribbon obstruction which arises in this setting.

Joint work with Patricia Cahn.

Sudipta Kolay (Georgia Tech)

Braided Embeddings

The theory of braids has been very useful in the study of classical knot theory. One can hope that higher dimensional braids will play a similar role in higher dimensional knot theory. In this talk we will introduce the concept of braided embeddings of manifolds, and discuss existence, lifting and isotopy problems for braided embeddings.

Christian Lange (Univ. Köln)

Orbifolds from a metric viewpoint

In the talk we explain the concept of an orbifold and its covering space theory from a metric viewpoint. The presence of a metric makes the story less technical compared to the smooth setting. We discuss some geometric and topological questions and results related to orbifolds.

Livio Liechti (Univ. Paris 6)

A graph model for positive braid links

We discuss a model for positive braid links, given by certain graphs which are embedded in the plane and carry an acyclic orientation. By a result of Stallings, the exterior of a positive braid link has the structure of a surface bundle over the circle, where the monodromy is a product of positive Dehn twists along simple closed curves. We discuss a graph model for positive braid links, using the key observation that such a surface bundle is determined by how the curves along which we twist intersect and in which order they get twisted along.

This is joint work with Sebastian Baader and Lukas Lewark.

Participants & Programme

Jeffrey Meier (Univ. Georgia)

Fibered homotopy-ribbon knots, the Generalized Property R Conjecture, and trisections

I'll describe recent work (joint with Alex Zupan) that draws connections between the study of fibered homotopy-ribbon knots in the three-sphere with the study of trisections of homotopy four-spheres, giving a new approach to the Generalized Property R Conjecture along the way. One application is the classification of fibered homotopy-ribbon disks bounded by square knots.

Jun Ueki (Univ. Tokyo)

The profinite completions of knot groups and twisted Alexander polynomials

In several occasions we can distinguish two knots by comparing the homology torsions of their finite covers. This leads us to the following question: What topological information does the profinite completions of knot groups know? In my talk, we study profinite rigidity of twisted Alexander polynomials of knots associated to representations over the ring O of S -integers of a number field. We prove it under some mild conditions, and also present some inevitable finite ambiguity, by investigating the completed group ring $\widehat{O}[[t^{\mathbb{Z}}]]$ and some Iwasawa modules.

Anderson Vera (Univ. Strasbourg)

Johnson-Levine homomorphisms and the tree reduction of the LMO functor

The interaction between the study of 3-manifolds and that of the mapping class group (MCG) is well known. In some sense, the algebraic structure of the MCG and of its subgroups is reflected in the topology of 3-manifolds. For instance, the subgroup of homeomorphisms acting trivially in homology, known as the Torelli group, is tied to homology 3-spheres. For this family of 3-manifolds there exists a powerful (quantum) invariant called the LMO invariant, which is a further generalization of the Casson invariant and that is quite mysterious, in part because its construction is very indirect. This invariant can be extended to a topological quantum field theory from a category of cobordisms to a category of univalent graphs, called the LMO functor. In this talk we show that for a special kind of cobordisms, a part of the tree-like graphs in the image by the LMO functor can be interpreted as the diagrammatic version of the Johnson-Levine homomorphisms (homomorphisms similar to the classical Johnson homomorphisms).

Stefan Witzel (Univ. Bielefeld)

The complex of boundary braids

A boundary braid in a solid cylinder is defined by the condition that certain strands stay in the boundary cylinder. I will present decomposition results for boundary braids. Specifically, the dual braid complex of braids with n strands of which k are boundary parallel decomposes as a metric direct product into the dual braid complex of braids with $n-k$ strands and a Euclidean polyhedron. Conjecturally, the dual braid complex is $CAT(0)$ and therefore the braid group is a $CAT(0)$ group. By our result the dual braid complex on n strands is covered by n subcomplexes which would be $CAT(0)$ in an inductive proof. This is joint work with Jon McCammond and Michael Dougherty.