



Winter Braids Lecture Notes

Arunima Ray

Slice knots and knot concordance

Vol. 8 (2021), Course n° II, p. 1-31.

<https://doi.org/10.5802/wbln.39>



Texte mis en ligne dans le cadre du
Centre Mersenne pour l'édition scientifique ouverte
<http://www.centre-mersenne.org/>
e-ISSN : 2426-0312

Slice knots and knot concordance

ARUNIMA RAY

Abstract

These notes were prepared to accompany a sequence of three lectures at the conference Winterbraids XI in Dijon, held in December 2021. In them, we provide an introduction to slice knots and the equivalence relation of concordance. We explain some connections between slice knots and exotic smooth structures on \mathbb{R}^4 . We also introduce filtrations of the knot concordance groups and satellite operations.

Overview

Slice knots were first defined in 1958 by Fox and Milnor and have since become a flourishing field of study. While they were originally considered in the context of resolving singularities of surfaces in 4-manifolds, numerous other connections to questions in 3- and 4-manifold topology have been discovered. A notable highlight: every knot which is topologically slice but not smoothly slice gives rise to an exotic smooth structure on \mathbb{R}^4 ; more on this in Section 2.

The goal of these lecture notes is to provide an overview of the basic notions in the study of slice knots and knot concordance, leading up to a small selection of recent developments. Due to the limitations of time we will barely scratch the surface of this active and vibrant area. Nevertheless we will attempt to provide some references and pointers to other resources.

Exercises

There are exercises throughout the lecture notes, which are also compiled in a list at the end. The problems are separated into three levels. **Green** \triangle exercises are usually straightforward and should be attempted if you are seeing this material for the first time. Prerequisites are courses in introductory geometric and algebraic topology. **Orange** \square exercises are for readers who are already comfortable with some of the terminology; they may require nontrivial input from outside these lectures, which we have tried to indicate as hints. Finally, **red** \circ exercises are challenge problems. Open problems will be marked as such, and do not intersect with the exercises.

Relationship to lectures

Some details and exercises in these notes were not mentioned in the accompanying lectures. The order of topics has also been slightly modified. Many of the exercises are new.

Conventions

Homeomorphism of manifolds is denoted by the symbol \approx , while diffeomorphism is denoted by \cong . All manifolds are assumed to be oriented. All knots are assumed to be tame and oriented, i.e. (images of) smooth embeddings $S^1 \hookrightarrow S^3$, where both the domain and codomain are oriented by hypothesis. The set of nonnegative integers is denoted by \mathbb{N} .

Acknowledgements

I am grateful to the organisers of Winterbraids XI in Dijon for an excellent conference, especially considering the Covid-19 pandemic, and to the attendees for their lively participation and many questions. Many thanks are due as well to the anonymous referee for their helpful comments.

1. Slice and ribbon knots

Recall that a knot $K: S^1 \hookrightarrow S^3$ is *trivial* or the *unknot* if and only if it bounds an embedded disc in S^3 . We will often conflate a knot and its image, and it will be important that both S^1 and S^3 are oriented. See Figure 1.1 for some examples of knots. Here, by definition, a knot is trivial if it can be deformed via an ambient isotopy to the round unit circle $S^1 \subseteq \mathbb{R}^3 \subseteq S^3$. For more on classical knot theory, see e.g. [Fox62, Rol90, Liv93].




Figure 1.1: The two knots on the left are trivial. The three knots on the right are not trivial, and are respectively called the *right-handed trefoil*, the *left-handed trefoil*, and the *figure eight* knot.

Slice knots are, in some sense, generalised trivial knots – we still ask for the knot to bound an embedded disc, but it has the freedom of an additional (half-)dimension.

Definition 1.1. A knot $K \subseteq S^3 = \partial B^4$ is *smoothly slice* if it bounds a smoothly embedded disc, called a *smooth slice disc*, in B^4 .¹

We begin by discussing a number of elementary ways to construct new slice knots from a given slice knot. Let $K: S^1 \hookrightarrow S^3$ be a knot. The knot obtained by reversing the orientation of S^1 is called the *reverse* of K , denoted by rK . The knot obtained by reversing the orientation of S^3 is called the *mirror image* of K , denoted by \bar{K} . For example, the left-handed trefoil is the mirror image of the right-handed trefoil. Both trefoils are isotopic to their reverses. The figure eight is isotopic to both its reverse and its mirror image.

Proposition 1.2 (Exercise Δ 3). Let $K \subseteq S^3$ be a knot. Show that K is smoothly slice if and only if rK is smoothly slice if and only if \bar{K} is smoothly slice.

Given knots $K, J \subseteq S^3$, we can form their *connected sum* by taking the (oriented!) connected sum of pairs $(S^3, K) \# (S^3, J)$; see Figure 1.2. Given diagrams of knots K and J we can form a diagram of the connected sum by connecting the two diagrams by a trivial band, as shown

¹In 1958, Fox and Milnor called these *null-equivalent* knots, in unpublished work. Fox adopted the term *slice knot*, proposed by Edward Moise, in [Fox62].

in the figure, taking care to match the original orientations. This is a special case of a *band sum* defined later in this section.




Figure 1.2: The connected sum of the figure eight and the right-handed trefoil from Figure 1.1.

Proposition 1.3 (Exercise Δ 4). *If the knots $K, J \subseteq S^3$ are smoothly slice, then so is $K \# J$.*

By ambient Morse theory, we can isotope any smooth slice disc so that the level sets with respect to the radial function on B^4 are links, i.e. (the image of) a smooth embedding of a disjoint union of circles in S^3 , except at finitely many radii where the level sets have singularities. Since we are considering a disc in a 4-manifold, there are only three possible types of singularities: minima, saddles, and maxima. See Figure 1.3 for a picture and [Mil65; Mil63; Mat02; Nic11; GS99, Sections 4.2 and 6.2] for more on Morse theory.




Figure 1.3: The axes schematically indicate B^4 , where the coordinates x, y, z describe $\mathbb{R}^3 \subseteq S^3$ and the coordinate $w \geq 0$ indicates depth, so that S^3 occurs at $w = 0$. In both cases we have a knot in $\mathbb{R}^3 \subseteq S^3 = \partial B^4$. On the left we have a trivial knot, as detected by the embedded disc in S^3 shaded grey. On the right we have a smoothly slice knot, with a smooth slice disc lying in B^4 . The disc is in a particularly nice position with respect to the radial function on B^4 . At most w -slices, the disc appears as a link, i.e. an embedding of a disjoint union of circles. At finitely many values of w , we see singularities. There are three types of singularities: minima (red), saddles (orange), and maxima (blue). See Figure 1.4 for a more precise diagram.

Smooth slice discs, once they have been perturbed to respect the radial Morse function as above, can be described in terms of movies. More specifically, we consider the radial level sets as the radius increases from 0 to 1, with the latter corresponding to $\partial B^4 = S^3$ containing a smoothly slice knot. At each minimum, an unknot is born, split from every other component, if any. At every saddle singularity, two portions of a link either merge together, or split apart. At a maximum, an unknotted component, split from everything else, disappears. See Figure 1.4 for an example.

We especially like smooth slice discs with only two types of singularities, as in the following definition.




Figure 1.4: Describing a ribbon disc as a movie. At the very bottom, we have two minima. At a slightly larger radius in B^4 , the minima give rise to (split) unknots, i.e. an unlink. The subsequent cross sections, until the orange saddle singularity, describe an isotopy of this unlink. At the saddle singularity, two portions of the previous unlink touch each other, and at a slightly higher radius we see a ribbon knot.

Definition 1.4. A knot $K \subseteq S^3 = \partial B^4$ is ribbon if it bounds a smoothly embedded disc in B^4 with only minima and saddles. Such a disc is called a *ribbon* disc.

Given a link $L \subseteq S^3$, a band B is a copy of $[0, 1] \times [0, 1]$ embedded in S^3 , with $\{0, 1\} \times [0, 1]$ lying on L (matching the orientations), and otherwise disjoint from L . The arc $[0, 1] \times \{1/2\}$ is called the *core* of B . A *band sum* of L along the band B is the result of removing $\{0, 1\} \times [0, 1]$ from L and gluing in $[0, 1] \times \{0, 1\}$; see Figure 1.5. When the components of $\{0, 1\} \times [0, 1]$ lie on distinct components of L the result is also called a *fusion* of L along B . Note that the core $[0, 1] \times \{1/2\}$ of a band can be an interesting arc in $S^3 \setminus L$, and in particular can link with the components of L . A band sum on L along B can be reversed by performing a dual band move, and corresponds to removing $[0, 1] \times \{0, 1\}$ from the band sum and gluing $\{0, 1\} \times [0, 1]$ back in. One notices that the result is isotopic to the original link L .




Figure 1.5: (a) A link L (black) and a band B (dark green). The core of B is shown in light green. (b) The result of the band sum on L along B is shown in black. The arc guiding the dual band move reversing the previous band move is shown in blue.

From the movie perspective, reading the ribbon disc from the bottom up, we see that any ribbon knot is a fusion of some trivial link. (Why is it a fusion, rather than an arbitrary band sum?) In other words, it can be obtained from an n -component trivial link for some n (corresponding to minima) and a collection of $n - 1$ bands (corresponding to saddles), by performing band sums, each of which joins two distinct components of the original link, and no two of which join the same pair of components. Alternatively, we could run the movie backwards. From this perspective, a ribbon knot is a knot where one can add $n - 1$ bands for some n (corresponding to the dual band moves described above), so that the result is an n -component unlink.

The following shows that ribbon knots can be described purely 3-dimensionally. This gives a useful method to detect if a given knot is ribbon.

Proposition 1.5 (Exercise □ 20). *A knot $K \subseteq S^3$ is ribbon if and only if it bounds a disc in S^3 with only ribbon singularities, i.e. singularities of the form shown in Figure 1.6.²*




Figure 1.6: Left: A *ribbon singularity* (shown in blue) of a disc (shown in grey) in S^3 . Right: The smaller disc region encircled in red on the horizontal sheet could be pushed radially into B^4 to resolve the singularity.

Proposition 1.6 (Exercise Δ 6). *For any knot $K \subseteq S^3$, the knot $K \# r\bar{K}$ is ribbon.*

From Definition 1.4, one sees directly that any ribbon knot is smoothly slice. (For the equivalent definition given in Proposition 1.5, one could push a small region of the ribbon disc in S^3 into B^4 to produce an embedded disc, as indicated in Figure 1.6.) The converse is an important open problem.

Conjecture 1.7 (Open, slice-ribbon conjecture, Fox [Fox61, Problem 25]). *Every smoothly slice knot is ribbon.*

²Such singular discs in S^3 are also sometimes called ribbon discs.

This conjecture has been established for certain families of knots, e.g. 2-bridge knots [Lis07] and most 3-strand pretzel knots [GJ11, Lec15], but there is no general approach to attacking it at the moment. Other work on the slice-ribbon conjecture using a similar strategy includes [Lec12, Bry17, Lon14, AKPR21]. Some potential counterexamples have been proposed, e.g. [GST10, AT16]. A very promising candidate had been the $(2, 1)$ -cable of the figure eight, which was known to be nonribbon for twenty years [Miy94] but whose sliceness status was unknown. It was recently shown to not be smoothly slice [DKM⁺22] (see also [ACM⁺23]). Notably it is still open whether it is topologically slice (Definition 1.10).

The subtlety of the conjecture is seen by the following proposition. Specifically, to show that a smoothly slice knot is ribbon it is not a good strategy to prove that a given smooth slice disc is ribbon.

Proposition 1.8 (Exercise \square 22). *There exist smooth slice discs that are not ambiently isotopic (relative to the boundary) to any ribbon disc.*

1.1. Topologically slice knots

So far we have worked strictly in the smooth setting. We can however loosen this restriction slightly. We will need the following definition.

Definition 1.9. An embedding $\varphi: (F, \partial F) \hookrightarrow (M, \partial M)$, i.e. a continuous map which is a homeomorphism onto its image, of a surface F in a 4-manifold M is said to be *locally flat* if for all $x \in F$ there is a neighbourhood U of $\varphi(x)$ such that $(U, U \cap \varphi(F))$ is homeomorphic to either $(\mathbb{R}^4, \mathbb{R}^2)$, in the case that $x \in \text{Int} F$, or to $(\mathbb{R}_+^4, \mathbb{R}_+^2)$, in the case that $x \in \partial F$.

The above gives rise to another notion of sliceness for knots.

Definition 1.10. A knot $K \subseteq S^3 = \partial B^4$ is *topologically slice* if it bounds a locally flat embedded disc, called a *topological slice disc*, in B^4 .

It is straightforward to see that any smoothly slice knot is topologically slice. Via a deep result of Quinn [Qui82, Theorem 2.5.1; FQ90, Theorem 9.3], a topological slice disc admits a tubular neighbourhood, i.e. we could equivalently define a knot K to be topologically slice if it bounds an embedded disc Δ in B^4 , admitting a neighbourhood homeomorphic to $\Delta \times D^2$, intersecting S^3 in a tubular neighbourhood of K .

Remark 1.11. We motivated slice knots by saying that they generalise trivial knots. Certainly the trivial knot is topologically slice. We should however be careful not to be too general. Given any knot $K \subseteq S^3$, the coned disc $\text{cone}(K) \subseteq \text{cone}(S^3) = B^4$ is an embedded disc bounded by K in B^4 . See Figure 1.7. As shown in the following proposition, this disc is not in general locally flat.

Proposition 1.12 (Exercise \triangle 7). *Let $K \subseteq S^3$ be a knot. The coned disc $\text{cone}(K) \subseteq \text{cone}(S^3) = B^4$ is locally flat if and only if K is the trivial knot.*

The following result of Freedman and Quinn gives a powerful method to construct topologically slice knots. For the definition of the Alexander polynomial, see e.g. [Rol90, Chapter 8].

Theorem 1.13 ([FQ90, Theorem 11.7B; GT04]). *Let $K \subseteq S^3$ be a knot. If the Alexander polynomial $\Delta_K(t) \doteq 1$, then K is topologically slice.*

As a consequence of the above theorem, the (untwisted) Whitehead double of any knot (Figure 1.8) is topologically slice, since it has Alexander polynomial one. On the other hand, many such knots are not smoothly slice.

Theorem 1.14. *There exist knots that are topologically slice but not smoothly slice, e.g. the positive clasped untwisted Whitehead double of the right-handed trefoil knot.*




Figure 1.7: Let $K \subseteq S^3$ be a knot. Thinking of B^4 as the cone on S^3 , we get an embedded disc $\text{cone}(K) \subseteq B^4$.




Figure 1.8: Whitehead doubling. On the left we have a knot Wh^+ in a solid torus $S^1 \times D^2$. In the middle we show the figure eight knot. Given an arbitrary knot K , the (positive clasped, untwisted) *Whitehead double* of K , denoted by $\text{Wh}^+(K)$, is the image of Wh^+ under a map identifying $S^1 \times D^2$ with a tubular neighbourhood of K , so that the longitude $S^1 \times \{*\}$ of $S^1 \times D^2$ is sent to the Seifert longitude of K . On the right we show the (positive clasped, untwisted) Whitehead double of the figure eight. The negative clasped Whitehead double $\text{Wh}^-(K)$ is defined in the same way, using as input the knot Wh^- in $S^1 \times D^2$ obtained from Wh^+ by changing both crossings. Whitehead doubling is a special case of the *satellite construction* on knots (Definition 5.1).

The above was first shown by Casson and Akbulut, in independent unpublished work, using Donaldson's diagonalisation theorem [Don83, Don86, Don87]. The first published accounts are by Gompf [Gom86], Cochran–Lickorish [CL86], and Cochran–Gompf [CG88].

However, for many examples of Whitehead doubles, it is still open whether they are smoothly slice. For the next question, let L denote the left-handed trefoil and let 4_1 denote the figure eight.

Question 1.15 (Open). Is $\text{Wh}^+(L)$ or $\text{Wh}^\pm(4_1)$ smoothly slice?

It is not difficult to see that if the knot K is smoothly slice, then so is $\text{Wh}^\pm(K)$; see [Exercise \$\triangle\$ 8](#). The converse is an interesting open question, generalising Question 1.15.

Conjecture 1.16 (Open [Kir97, Problem 1.38]). *Let $K \subseteq S^3$ be a knot. The knot $\text{Wh}^\pm(K)$ is smoothly slice if and only if K is smoothly slice.*

As mentioned earlier, Donaldson's diagonalisation theorem was used to show that the positive clasped untwisted Whitehead double of the right-handed trefoil is not smoothly slice. This is a relatively subtle sliceness obstruction, since e.g. it does not obstruct topological

sliceness. There are numerous other slicing obstructions, including many topological sliceness obstructions. We will return to this topic in Section 4, where we will explain a method to arrange sliceness obstructions in order of strength, in some sense. We defer that discussion to first describe some connections between slice knots and 4-manifold topology in the upcoming section.

1.2. Slice genus

We have seen that not every knot $K \subseteq S^3$ bounds a (smooth or topological) slice disc in B^4 . However, since $H_1(B^4; \mathbb{Z}) = 0$, we know that every knot bounds at least some embedded smooth or locally flat connected surface in B^4 . This gives rise to the following notion.

Definition 1.17. Let $K \subseteq S^3 = \partial B^4$ be a knot. The *smooth slice genus* of K is the minimum genus of a compact, connected, oriented surface $\Sigma \subseteq B^4$ such that $\partial \Sigma = K$. The *topological slice genus* of K is defined analogously, except that Σ is only required to be locally flat.

The slice genus of a knot gives a natural measure of how far it is from being slice, in either the smooth or topological settings. Many obstructions to sliceness, including many coming from classical invariants like signature or from modern technology like Heegaard-Floer homology or Khovanov homology, in fact provide lower bounds for the slice genus. See [Fel16] for a generalisation of Theorem 1.13 to the topological slice genus. Unfortunately for lack of time we will not discuss the slice genus further in these notes, except for in some questions in Section 4.

1.3. Links

The notions in this section can be generalised to links, i.e. embeddings $\sqcup S^1 \hookrightarrow S^3$. For example, we have the following definition.

Definition 1.18. A link $L \subseteq S^3 = \partial B^4$ is said to be (*strongly*) *smoothly slice* if the components bound a collection of pairwise disjoint smooth slice discs in B^4 . If the discs are only locally flat, the link is said to be (*strongly*) *topologically slice*.

There is a parallel notion of *weak* sliceness of links, in both the topological and smooth settings. For this we do not require a collection of pairwise disjoint embedded discs, but rather just some embedded planar surface (either smooth or locally flat).

Unfortunately we will not have time to pursue links thoroughly. We limit ourselves to saying that the open questions mentioned in this section have analogues for links: it is open whether the slice-ribbon conjecture is true for links and whether the Whitehead double of a link L is smoothly slice if and only if L is smoothly slice. It is also open whether the Whitehead double of every link is topologically slice, and an answer would have deep consequences for fundamental open questions in 4-manifold topology (see e.g. [KOPR21]).

2. Exotic smooth structures on \mathbb{R}^4

By definition, a smooth structure on a topological manifold M is a maximal atlas of charts on M such that the transition maps are smooth. *A priori* a given topological manifold can admit finitely many (including none) or infinitely many smooth structures. The study of existence and uniqueness questions for smooth structures comprises the field of *smoothing theory*. Notably, the sphere S^7 has 28 distinct smooth structures up to orientation preserving diffeomorphism [Mil56, KM63], and whether S^4 has more than one smooth structure is the content of the (still open) smooth 4-dimensional Poincaré conjecture. Smoothing theory is well developed in dimensions five and higher, and for example, using technology of Kirby-Siebenmann from [KS77], it can be shown that closed manifolds of dimension five and higher have finitely many (possibly zero) smooth structures. Smoothing closed 4-manifolds is more

complicated. There exist topological 4-manifolds which do not admit any smooth structures, such as Freedman’s $E8$ -manifold [Fre82], while there are closed 4-manifolds admitting infinitely many smooth structures, such as the $K3$ surface [FS98].

The existence and uniqueness questions for smooth structures can also be asked for non-compact manifolds. The following result shows that Euclidean spaces are well behaved assuming $n \neq 4$.

Theorem 2.1 ([Moi77, Sta62]). *For $n \neq 4$ there is a unique smooth structure on \mathbb{R}^n , up to diffeomorphism.*

By contrast, there are *uncountably* many smooth structures on \mathbb{R}^4 [Tau87].

Definition 2.2. Let $\mathbb{R}_{\text{std}}^4$ denote \mathbb{R}^4 with its standard smooth structure. A smooth 4-manifold which is homeomorphic to $\mathbb{R}_{\text{std}}^4$, but not necessarily diffeomorphic to $\mathbb{R}_{\text{std}}^4$, is called an \mathbb{R}^4 -homeomorph. An \mathbb{R}^4 -homeomorph which is not diffeomorphic to $\mathbb{R}_{\text{std}}^4$ is called an *exotic* \mathbb{R}^4 , and such a smooth structure is called an *exotic* smooth structure on \mathbb{R}^4 .

There are two distinct types of \mathbb{R}^4 -homeomorphs, as per the following definition.

Definition 2.3. Let \mathcal{R} be an \mathbb{R}^4 -homeomorph. The manifold \mathcal{R} is said to be *small* if it admits a smooth embedding $\mathcal{R} \hookrightarrow \mathbb{R}_{\text{std}}^4$. It is called *large* otherwise.³

Knots and links feature prominently in the known constructions of exotic \mathbb{R}^4 s. In this section we describe two such constructions, one of large exotic \mathbb{R}^4 s, and one of small exotic \mathbb{R}^4 s.

2.1. Large exotic \mathbb{R}^4 s using 0-traces

For any knot K we have the *0-trace*, which is by definition the smooth 4-manifold obtained by adding a 0-framed 2-handle to B^4 along $K \subseteq S^3$, and then smoothing corners. One of the two key ingredients in the forthcoming construction is the following characterisation of slice knots.

Lemma 2.4 (Trace embedding lemma, Exercise □ 23). *Let $K \subseteq S^3$ be a knot. The 0-trace $X_0(K)$ admits a smooth (resp. locally collared) embedding into $\mathbb{R}_{\text{std}}^4$ if and only if K is smoothly (resp. topologically) slice.*

We will also need the following powerful result.

Theorem 2.5 ([Qui82, Corollary 2.2.3; FQ90, Theorem 8.2, Section 8.7]). *Let M be a connected, noncompact 4-manifold. If desired, fix a smooth structure on any collection of connected components of ∂M . There is a smooth structure on M extending the given smooth structure on (a subset of) ∂M .*

We are now ready for the construction.

Construction 2.6 ([Gom85, Lemma 1.1; GS99, Exercise 9.4.23]). *Given a knot $K \subseteq S^3$ which is topologically slice but not smoothly slice there exists \mathcal{R} , a large exotic \mathbb{R}^4 , with a smooth embedding $X_0(K) \hookrightarrow \mathcal{R}$.*

Proof. Let K be a knot which is topologically slice but not smoothly slice. Then there is a locally collared embedding $\varphi: X_0(K) \hookrightarrow \mathbb{R}^4$ by the trace embedding lemma. Then the image $\varphi(X_0(K))$ inherits a smooth structure from $X_0(K)$ since it is homeomorphic to the latter. Since φ is locally collared, we see that $\mathbb{R}^4 \setminus \text{Int} \varphi(X_0(K))$ is a manifold, and we can check using the Mayer–Vietoris sequence that it is connected. Since $X_0(K)$ is compact and \mathbb{R}^4 is not, we

³There are two slightly different definitions of small/large \mathbb{R}^4 -homeomorphs in the literature. Ours coincides with the definition of Scorpan’s book [Sco05]. In the book by Gompf–Stipsicz [GS99], a large \mathbb{R}^4 -homeomorph is one which contains a compact subset that does not embed in $\mathbb{R}_{\text{std}}^4$, and an \mathbb{R}^4 -homeomorph is called small if it is not large. It is not known whether the two definitions are equivalent.

also know that $\mathbb{R}^4 \setminus \text{Int} \varphi(X_0(K))$ is noncompact. So by Theorem 2.5, we extend the smooth structure on $\varphi(\partial X_0(K))$ to the rest of $\mathbb{R}^4 \setminus \text{Int} \varphi(X_0(K))$. This produces a smooth structure on \mathbb{R}^4 , and we denote the corresponding smooth 4-manifold by \mathcal{R} .

Suppose that $\mathcal{R} \cong \mathbb{R}_{\text{std}}^4$. Then by construction we have a smooth embedding $X_0(K) \hookrightarrow \mathcal{R} \cong \mathbb{R}_{\text{std}}^4$, so K is smoothly slice by the trace embedding lemma, which is a contradiction. Therefore \mathcal{R} is an exotic \mathbb{R}^4 .

Indeed, since $X_0(K) \subseteq \mathcal{R}$ has no smooth embedding in $\mathbb{R}_{\text{std}}^4$, we see that \mathcal{R} is large.⁴ \square

2.2. Small exotic \mathbb{R}^4 s using ribbon disc exteriors

The construction in this section will use *Casson handles*. These are smooth noncompact 4-manifolds, constructed by Casson in [Cas86] as approximations of (open) 2-handles. The boundary of a Casson handle C , called the *attaching region* ∂C , is identified with the open solid torus $S^1 \times \text{Int} D^2$, and from the Kirby diagrams of Casson handles (see, e.g. [Kir89, Chapter XII; GS99, Chapter 6; Fre82, Section 2]), one observes that every Casson handle C admits a smooth embedding $(C, \partial C) \hookrightarrow (D^2 \times D^2, S^1 \times \text{Int} D^2)$, extending the aforementioned identification on the boundary. In [Fre82], Freedman showed that any Casson handle C is homeomorphic, relative to its attaching region, to an open 2-handle, i.e. $(C, \partial C) \approx (D^2 \times \text{Int} D^2, S^1 \times \text{Int} D^2)$, again extending the aforementioned identification on the boundary.

Construction 2.7. Given a smoothly slice link L with n components, and a collection of Casson handles $\{C_1, \dots, C_n\}$, there exists a small \mathbb{R}^4 -homeomorph \mathcal{R}_L .

Proof. Let $\{\Delta_i\}$ be a collection of smooth slice discs for L . Consider the complement $B^4 \setminus \bigcup \nu \Delta_i$, of open tubular neighbourhoods of the discs. If we glue in 2-handles along the meridians of the components of L , we will get back B^4 . However, we could instead glue in the Casson handles $\{C_1, \dots, C_n\}$ along those meridians. Let \mathcal{R}_L denote the result of gluing in Casson handles to $B^4 \setminus \bigcup \nu \Delta_i$ along the meridians of L , and then removing all the remaining boundary. Since Casson handles are homeomorphic to open 2-handles relative to the attaching region we see that \mathcal{R}_L is an \mathbb{R}^4 -homeomorph. Since every Casson handle embeds in a standard 2-handle, respecting the attaching region, we also see that \mathcal{R}_L admits a smooth embedding into $\mathbb{R}_{\text{std}}^4$, and so by definition \mathcal{R}_L is small. \square

For certain choices of L and Casson handles $\{C_1, \dots, C_n\}$, it can be shown that \mathcal{R}_L is not diffeomorphic to $\mathbb{R}_{\text{std}}^4$. For examples of this, see [DMF92, BG96]. Roughly speaking, showing that such an \mathcal{R}_L is exotic involves embedding it appropriately within a simply connected h -cobordism which is known to not be a smooth product. The known examples use ribbon links and are therefore called *ribbon \mathbb{R}^4 s*. All known small exotic \mathbb{R}^4 s are ribbon \mathbb{R}^4 s. The simplest known exotic ribbon \mathbb{R}^4 is built using the complement of a standard ribbon disc for the 9_{46} knot (also called the $(3, -3, 3)$ pretzel knot), and the Casson handle built using self-plumbed 2-handles within a single, positive self-plumbing at each stage [BG96] – this explicit construction yields a description of the corresponding exotic \mathbb{R}^4 as the interior of an infinite but rather simple handlebody.

2.3. Universal exotic \mathbb{R}^4

Finally, there is the Freedman–Taylor universal exotic \mathbb{R}^4 [FT86], denoted by \mathcal{U} , which has the remarkable property that every \mathbb{R}^4 -homeomorph \mathcal{R} admits a smooth embedding $\mathcal{R} \hookrightarrow \mathcal{U}$. The construction of \mathcal{U} also involves solving slicing problems for knots and links (via Kirby diagrams for Casson handles) and applying Quinn’s result (Theorem 2.5). Unfortunately the construction is beyond the scope of the lectures and we invite the reader to learn more about the construction in [FT86] on their own.

⁴This holds for either definition of “large”.

3. The knot concordance groups

So far we have considered sliceness as a generalisation of triviality for knots. Similarly, there is a generalisation of isotopy, called *concordance*. Specifically, we have the following definition.

Definition 3.1. Knots $K, J \subseteq S^3$ are said to be smoothly (resp. topologically) *concordant* if they cobound a smooth (resp. locally flat) embedded annulus in $S^3 \times [0, 1]$; see Figure 3.1. More specifically, we consider $K \subseteq S^3$ as lying in $S^3 \times \{1\}$ and $J \subseteq S^3$ as lying in $S^3 \times \{0\}$, and assert the existence of a *concordance* $A = S^1 \times [0, 1] \hookrightarrow S^3 \times [0, 1]$, with $S^1 \times \{0\}$ mapping to J and $S^1 \times \{1\}$ mapping to K , where the embedding of A is either smooth or locally flat, as needed.




Figure 3.1: A schematic picture showing a concordance between two knots $J \subseteq S^3 \times \{0\}$ and $K \subseteq S^3 \times \{1\}$.

Proposition 3.2 (Exercise \triangle 9). *A knot $K \subseteq S^3$ is smoothly (resp. topologically) slice if and only if it is smoothly (resp. topologically) concordant to the unknot.*

Recall that by our conventions, all the manifolds in the definition of concordance are oriented. Using the standard orientation of $[0, 1]$, we notice that $S^3 \times \{1\}$ inherits the standard orientation of S^3 while $S^3 \times \{0\}$ inherits the opposite orientation. Similarly a concordance (i.e. a copy of $S^1 \times [0, 1]$) induces opposite orientations on its boundary components. Under these orientations, we see that if K and J are concordant knots, then the claimed annulus joins $K \subseteq S^3 \times \{1\} = S^3$ to $rJ \subseteq S^3 \times \{0\} = -S^3$. The latter knot is by definition $\overline{rJ} = r\bar{J}$.

When speaking about slice knots we often ignore these orientation conventions, but they are key to proving the following fact.

Proposition 3.3 (Exercise \square 24). *Two knots $K, J \subseteq S^3$ are smoothly (resp. topologically) concordant if and only if $K \# r\bar{J}$ is smoothly (resp. topologically) slice.*

Similar to isotopy, concordance (both smooth and topological) is an equivalence relation. However, unlike isotopy (Exercise \square 25), the equivalence classes under concordance form a group under the connected sum operation.

Proposition 3.4 (Exercise \triangle 10). *Smooth (resp. topological) concordance is an equivalence relation on knots.*

Definition 3.5 (Concordance groups). The *smooth concordance group*, denoted by $\mathcal{C}^{\text{diff}}$, consists of smooth concordance classes of knots, under the operation of connected sum.

Similarly, the *topological concordance group*, denoted by \mathcal{C}^{top} , consists of topological concordance classes of knots, under the operation of connected sum.

We often conflate a knot K with its concordance class $[K]$. We asserted but did not prove that concordance classes form a group. This is the combination of the following proposition and [Exercise \$\Delta\$ 2](#).

Proposition 3.6 ([Exercise \$\Delta\$ 11](#)). *Let $K \subseteq S^3$ be a knot. In either $\mathcal{C}^{\text{diff}}$ or \mathcal{C}^{top} , the inverse of $[K]$ is the class of $-K := r\bar{K}$. The identity element in $\mathcal{C}^{\text{diff}}$ (resp. \mathcal{C}^{top}) is the class of the unknot, equivalently the class of any smoothly (resp. topologically) slice knot.*

By [Exercise \$\Delta\$ 2](#), the concordance groups are abelian, so it is tempting to speculate that they cannot be particularly complicated. This is somewhat of a red herring. There is an algebraic version of the knot concordance group, called the *algebraic concordance group*, denoted by \mathcal{AC} . This group has a couple of equivalent definitions, one in terms of Seifert matrices and another in terms of Alexander modules and Blanchfield forms – see [Lev69a, Lev69b, Kea73, Tro73, Kea75a, Kea75b, Sto77, Tro78] for further details. Given the definitions, it is not hard to see that there is a surjection

$$\Phi: \mathcal{C}^{\text{top}} \twoheadrightarrow \mathcal{AC},$$

given by taking a knot to the class of its Seifert matrix, or of its Alexander module and Blanchfield form. It was shown by J. Levine [Lev69b] (see also Stoltzfus [Sto77]) that

$$\mathcal{AC} \cong \mathbb{Z}^\infty \oplus \mathbb{Z}/2^\infty \oplus \mathbb{Z}/4^\infty,$$

so \mathcal{C}^{top} is quite large.

Notice that \mathcal{AC} has many elements of order two. So do $\mathcal{C}^{\text{diff}}$ and \mathcal{C}^{top} by the following proposition.

Proposition 3.7 ([Exercise \$\Delta\$ 12](#)). *A knot $K \subseteq S^3$ is said to be *negative amphichiral* if it is isotopic to $r\bar{K}$. Every negative amphichiral knot has order at most two in $\mathcal{C}^{\text{diff}}$ and \mathcal{C}^{top} . As a result, the figure eight knot has order two in $\mathcal{C}^{\text{diff}}$ and \mathcal{C}^{top} .*

Conjecture 3.8 (Open, Gordon [Kir97, Problem 1.94]). *Let $K \subseteq S^3$ be a knot. If $K\#K$ is slice, then K is smoothly concordant to a negative amphichiral knot.*

By contrast, there are no known candidates for knots of order four (note that \mathcal{AC} has many such elements). We have the following open question.

Question 3.9 (Open, see [Kir97, Problem 1.32]). *Does the map Φ split? In particular, does \mathcal{C}^{top} (or $\mathcal{C}^{\text{diff}}$) have any elements of order four? Indeed, are there any knots with finite order $\neq 0, 2$ in $\mathcal{C}^{\text{diff}}$ or \mathcal{C}^{top} ?*

The kernel of the map Φ has also been an object of much study, and is the focus of the next section.

Remark 3.10. One may also consider high-dimensional knots, i.e. embeddings $S^n \hookrightarrow S^{n+2}$ for arbitrary n . As in the classical case, one then has corresponding concordance and algebraic concordance groups. Indeed, for even n , every embedding $S^n \hookrightarrow S^{n+2}$ is slice, in the smooth, piecewise linear, or smooth categories [Ker65, Théorème III.6; Ker71, Theorem 1]. For odd $n \geq 5$, the topological and piecewise linear concordance groups of such embeddings is precisely in bijection with the corresponding algebraic concordance group (for $n = 3$ there is a discrepancy of $\mathbb{Z}/2$ related to Rochlin’s theorem) [Lev69b, Ker71]. In the smooth category, the odd high-dimensional concordance groups are also closely determined by the corresponding algebraic concordance group, with a discrepancy of \mathbb{Z}/k for some k – this discrepancy is related to the existence of exotic smooth structures on high-dimensional spheres and the famous Kervaire invariant problem (see [HHR16] for an account of the history of this problem). Roughly speaking, the map from knot concordance classes to the algebraic concordance group is still injective, as in the case for topological or piecewise linear concordance in high dimensions, but we no longer get a surjection since the construction in the realisation step may produce an exotic sphere rather than a standard one.

From the definitions of the concordance groups it should be clear that there is a surjection

$$(3.1) \quad \Psi: \mathcal{C}^{\text{diff}} \twoheadrightarrow \mathcal{C}^{\text{top}}.$$

As mentioned in Section 1, there exist knots that are topologically slice but not smoothly slice. These (or rather, their smooth concordance classes) lie in the kernel of Ψ .

Definition 3.11. The subset $\mathcal{T} \subseteq \mathcal{C}^{\text{diff}}$ consists of the smooth concordance classes of topologically slice knots.

The structure of \mathcal{T} is another key area of research (see e.g. [Gom86, CG88, End95, HK12, HLR12, CHH13, Hom14, CH15, Hom15, HKL16, OSS17, Hom19, CK21, DHST21, KL22]). One method of studying this structure is described in the following section.

4. Filtrations of the knot concordance groups

4.1. Overview

As mentioned in Remark 3.10, the high-dimensional codimension two concordance groups of odd-dimensional knots are controlled almost entirely by the corresponding algebraic analogues, and in almost all cases they are isomorphic. Therefore, it was an interesting question for several years whether the knot concordance groups \mathcal{C}^{top} and the algebraic knot concordance group \mathcal{AC} are isomorphic also in the classical dimension, in other words, whether the map Φ is a bijection. Since it is easily shown to be a surjection, the main question was whether Φ is injective. The first nontrivial elements in the kernel of Φ were given by Casson and Gordon in [CG78, CG86]. Then Cochran, Orr, and Teichner [COT03] defined the *solvable filtration* of $\mathcal{C}^{\text{diff}}$, denoted by $\{\mathcal{F}_i\}_{i \in \frac{1}{2}\mathbb{N}}$, satisfying

$$\bigcap_{i \in \frac{1}{2}\mathbb{N}} \mathcal{F}_i \subseteq \cdots \subseteq \mathcal{F}_{n+1} \subseteq \mathcal{F}_{n.5} \subseteq \mathcal{F}_n \subseteq \cdots \subseteq \mathcal{F}_0 \subseteq \mathcal{C}^{\text{diff}}.$$

Remark 4.1. Cochran–Orr–Teichner defined the filtration in the locally flat category, for the topological concordance group \mathcal{C}^{top} . In light of future developments we give the definition in the smooth category. For the relationship between the two notions see Proposition 4.7 and Corollary 4.8.

We will give the precise definition of the terms of this filtration presently. For now, we note that it is highly nontrivial (see Theorem 4.9), and it subsumes several other invariants, e.g. a knot K is in \mathcal{F}_0 if and only if $\text{Arf}(K) = 0$; it is in $\mathcal{F}_{0.5}$ if and only if it is algebraically slice; and if K is in $\mathcal{F}_{1.5}$ then all its Casson–Gordon sliceness obstructions vanish (see [Exercise 31](#)). We will also see in Theorem 4.12 that $\mathcal{T} \subseteq \bigcap_{i \in \frac{1}{2}\mathbb{N}} \mathcal{F}_i$.

Question 4.2 (Open). Is $\mathcal{T} = \bigcap_{i \in \frac{1}{2}\mathbb{N}} \mathcal{F}_i$?

On the other hand, as we mentioned in Section 1, we know that $\mathcal{T} \neq \{1\}$. By Theorem 4.12 the solvable filtration cannot effectively distinguish between smooth concordance classes of topologically slice knots. For this purpose, Cochran–Harvey–Horn [CHH13] defined the *bipolar filtration* of \mathcal{T} , denoted by $\{\mathcal{T}_n\}_{n \in \mathbb{N}}$, satisfying

$$\bigcap_{i \in \mathbb{N}} \mathcal{T}_i \subseteq \cdots \subseteq \mathcal{T}_{n+1} \subseteq \mathcal{T}_n \subseteq \cdots \subseteq \mathcal{T}_0 \subseteq \mathcal{T}.$$

We will define the bipolar filtration presently, and note here that it is also highly nontrivial (see Theorem 4.15).

Question 4.3 (Open). Is $\bigcap_{i \in \mathbb{N}} \mathcal{T}_i = \{1\}$?

4.2. Definitions of the filtrations

Both the solvable filtration of Cochran–Orr–Teichner and the bipolar filtration of Cochran–Harvey–Horn are motivated by the following characterisation of topologically slice knots. For a knot $K \subseteq S^3$, the 3-manifold obtained by performing 0-framed Dehn surgery on S^3 along K is denoted by $S_0^3(K)$.

Proposition 4.4. *Let $K \subseteq S^3$ be a knot. Then K is topologically slice if and only if $S_0^3(K) = \partial W$, where W is a compact, connected, oriented 4-manifold such that*

- (i) *inclusion induces an isomorphism $\mathbb{Z} \cong H_1(S_0^3(K); \mathbb{Z}) \rightarrow H_1(W; \mathbb{Z})$;*
- (ii) *$H_2(W; \mathbb{Z}) = 0$; and*
- (iii) *$\pi_1(W)$ is normally generated by the meridian $\mu_K \subseteq S_0^3(K)$.*

Proof. Assume that K is topologically slice, with a topological slice disc $\Delta \subseteq B^4$. As mentioned before, by work of Quinn [Qui82, Theorem 2.5.1; FQ90, Theorem 9.3] we know that Δ admits an open tubular neighbourhood $\nu\Delta \approx \Delta \times \mathring{D}^2$. Then let $W = B^4 \setminus \nu\Delta$. The boundary of W is a union of the knot exterior $S^3 \setminus \nu K$ and $\Delta \times S^1$, a solid torus. This shows that ∂W is the result of some Dehn surgery on S^3 along K . In order to see that we indeed get $S_0^3(K)$ as the boundary, note that the disc $\Delta \times \text{pt} \subseteq \Delta \times S^1$ is attached to a pushoff of K which has zero linking number with K , since in particular it bounds a disc in B^4 disjoint from $\Delta \times \{0\} \subseteq \nu\Delta$.

Then (iii) follows from the Seifert–van Kampen theorem since we obtain B^4 (which is simply connected) from W by gluing in a thickened 2-cell (i.e. the neighbourhood $\nu\Delta$) to W along $\mu_K \subseteq \partial W$.

To see (ii), consider the Mayer–Vietoris sequence for B^4 as the union $W \cup \nu\Delta = B^4$. Notice that $W \cap \nu\Delta \cong D^2 \times S^1$. Then we have

$$H_2(D^2 \times S^1; \mathbb{Z}) \rightarrow H_2(W; \mathbb{Z}) \oplus H_2(\nu\Delta; \mathbb{Z}) \rightarrow H_2(B^4; \mathbb{Z}),$$

so we see that $H_2(W; \mathbb{Z}) = 0$. From the same sequence, we also get that $H^2(W; \mathbb{Z}) \cong H^3(W; \mathbb{Z}) = 0$. Therefore, $H_1(W, \partial W; \mathbb{Z}) \cong H_2(W, \partial W; \mathbb{Z}) = 0$. Apply these to the long exact sequence for the pair $(W, \partial W)$, where we know that $\partial W \cong S_0^3(K)$:

$$H_2(W, S_0^3(K); \mathbb{Z}) \rightarrow H_1(S_0^3(K); \mathbb{Z}) \rightarrow H_1(W; \mathbb{Z}) \rightarrow H_1(W, S_0^3(K); \mathbb{Z})$$

This yields (i) and completes half of the proof.

For the converse direction, given W , we will glue in a 2-handle along $\mu_K \subseteq \partial W$, with a framing we describe presently, so that the result \mathcal{B} is homeomorphic to B^4 , within which we will locate a slice disc for the knot $K \subseteq \partial\mathcal{B}$. To do so, we will need to choose a framing of μ_K with the goal of recovering S^3 as $\partial\mathcal{B}$. We do this explicitly next. Experts can safely skip this level of detail, and instead work directly in a Dehn surgery diagram. We provide an explicit argument for the benefit of newcomers to the field.

This construction will use multiple copies of the 2-disc D^2 , which we indicate with subscripts, i.e. each D_i^2 below indicates a copy of D^2 . Use the Seifert longitude of K to identify a tubular neighbourhood $\nu K \subseteq S^3$ with $K \times D_0^2$, so the Seifert longitude is given by $K \times x$ for some $x \in \partial D_0^2$. By definition, we know that

$$S_0^3(K) = (S^3 \setminus (K \times D_0^2)) \cup_{\partial D_1^2 \times S^1} (D_1^2 \times S^1),$$

with the *surgery solid torus* $D_1^2 \times S^1$ attached to $K \times \partial D_0^2$ so that $\partial D_1^2 \times y$ for some $y \in S^1$ is identified with $K \times x$ and $z \times S^1$ for $z \in \partial D_1^2$ is identified with the meridian μ_K . Attach a 2-handle $D_1^2 \times D_2^2$ to W along the core of the surgery solid torus, i.e. along $0 \times S^1$, using the surgery solid torus as the requisite framing of a tubular neighbourhood of the core, and call the result \mathcal{B} . In other words we have:

$$\mathcal{B} := W \cup_{D_1^2 \times S^1} D_1^2 \times D_2^2,$$

Note that the core $0 \times S^1 \subseteq D_1^2 \times S^1$ is isotopic in $S_0^3(K)$ to $z \times S^1$, which we saw is identified with μ_K , so we are indeed attaching a 2-handle to W along μ_K as claimed.

Then $\partial\mathcal{B}$ is the result of surgery on $S_0^3(K)$ along $\mu(K)$. More precisely, we have

$$\begin{aligned} \partial\mathcal{B} &= (S_0^3(K) \setminus (D_1^2 \times S^1)) \cup_{S^1 \times \partial D_2^2} (S^1 \times D_2^2) \\ &= \left(((S^3 \setminus (K \times D_0^2)) \cup_{\partial D_1^2 \times S^1} (D_1^2 \times S^1)) \setminus (D_1^2 \times S^1) \right) \cup_{S^1 \times \partial D_2^2} (S^1 \times D_2^2) \\ &= (S^3 \setminus (K \times D_0^2)) \cup_{S^1 \times \partial D_2^2} (S^1 \times D_2^2), \end{aligned}$$

where we can check that $w \times \partial D_2^2$ for $w \in S^1$ is attached to the meridian μ_K . In other words, we have performed the ∞ -framed Dehn surgery on S^3 along K , producing $\partial\mathcal{B} = S^3$. In this copy of S^3 , we still have the knot K , represented as $\partial D_1^2 \times y$, so in particular the disc $D_1^2 \subseteq \mathcal{B}$ is a slice disc for K .

Moreover, by our hypotheses, we also see that \mathcal{B} is homotopy equivalent to B^4 . Then by the topological 4-dimensional Poincaré conjecture [Fre82, Theorem 1.6], we know that $\mathcal{B} \approx B^4$. This completes the proof. \square

We now give the definition of the solvable filtration. Recall that for a group G , the *ith derived subgroup*, denoted $G^{(i)}$, is inductively defined by setting $G^{(0)} = G$ and $G^{(i+1)} := [G^{(i)}, G^{(i)}]$ for each $i \geq 0$.

Definition 4.5 ([COT03, Definition 1.2]). Let $K \subseteq S^3$ be a knot and $n \in \mathbb{N}$. We say that K is (smoothly) *n-solvable* if $S_0^3(K) = \partial W$ where W is a smooth, compact, connected, oriented 4-manifold such that

- (i) the inclusion induces an isomorphism $\mathbb{Z} \cong H_1(S_0^3(K); \mathbb{Z}) \rightarrow H_1(W; \mathbb{Z})$; and
- (ii) $H_2(W; \mathbb{Z})$ has a basis consisting of smoothly embedded, closed, connected, oriented surfaces $\{L_i, D_i\}_{i=1}^k$, for some k , such that
 - (a) each L_i and D_i has trivial normal bundle, and
 - (b) the surfaces $\{L_i, D_i\}_i$ are pairwise disjoint, except that for each i , the surface L_i intersects D_i transversely once with positive sign; and
 - (c) for each i , we have $\pi_1(L_i) \subseteq \pi_1(W)^{(n)}$ and $\pi_1(D_i) \subseteq \pi_1(W)^{(n)}$, with respect to the inclusion induced maps.

The set of *n-solvable* knots is denoted by \mathcal{F}_n and the manifold W is called an *n-solution*.

We say that K is (smoothly) *n.5-solvable* if, in addition, for each i , we have $\pi_1(D_i) \subseteq \pi_1(W)^{(n+1)}$ with respect to the inclusion induced maps. The set of *n.5-solvable* knots is denoted by $\mathcal{F}_{n.5}$ and the manifold W is called an *n.5-solution*.

We leave it to the reader to show that, for each $n \in \frac{1}{2}\mathbb{N}$, the set \mathcal{F}_n is a subgroup of $\mathcal{C}^{\text{diff}}$ (Exercise $\triangle 13$). Note that condition (i) of Definition 4.5 and of Proposition 4.4 are the same. Condition (ii) in Definition 4.5 can be seen as a generalisation of condition (ii) in Proposition 4.4 – roughly speaking, if we assume that each surface L_i is a sphere, then one could do surgery on W along $\{L_i\}$, i.e. remove a tubular neighbourhood $L_i \times D^2$ and glue in $S^1 \times D^3$, for each i , and the resulting 4-manifold with boundary $S_0^3(K)$ would have trivial second homology. The condition that $\pi_1(L_i) \subseteq \pi_1(W)^{(n)}$ then measures how far away the homology class of L_i is from being represented by an immersed sphere: in the case that $\pi_1(L_i)$ is mapped to the trivial subgroup of $\pi_1(W)$, one could perform ambient surgery along half a symplectic basis of curves on L_i to transform it to an immersed sphere.

Remark 4.6. We could also add in Condition (iii) from Proposition 4.4 to Definition 4.5, without any loss to known results about the solvable filtration. *A priori* (but only conjecturally) that would lead to a different filtration.

In Definition 4.5, if the manifold W is not required to be smooth and the surfaces $\{L_i, D_i\}_i$ are only required to be locally flat embedded, we say that K is *topologically n -solvable* or *$n.5$ -solvable*. Denote the set of such knots by $\mathcal{F}_n^{\text{top}}$ and $\mathcal{F}_{n.5}^{\text{top}}$ respectively. *A priori* the filtrations $\{\mathcal{F}_n^{\text{top}}\}_{n \in \frac{1}{2}\mathbb{N}}$ and $\{\mathcal{F}_n\}_{n \in \frac{1}{2}\mathbb{N}}$ appear to be distinct, but indeed, they coincide, as mentioned in [CHL09, p. 1423; CHH13, p. 2105]. Since an explicit proof has not appeared in the literature before, we provide one next.

Proposition 4.7. *Let $K \subseteq S^3$ be a knot and $n \in \frac{1}{2}\mathbb{N}$. Then K is smoothly n -solvable if and only if it is topologically n -solvable.*

Proof. By forgetting smoothness, we see that if $K \in \mathcal{F}_n$ then $K \in \mathcal{F}_n^{\text{top}}$.

Assume that $K \in \mathcal{F}_n^{\text{top}}$ and let W denote a topological n -solution. First we note that the Kirby–Siebenmann invariant $\text{ks}(W) = 0$. To see this, observe that W is spin, since $H_1(W; \mathbb{Z}) \cong \mathbb{Z}$ has no 2-torsion and the intersection form is even. The Kirby–Siebenmann invariant of a compact, oriented, spin topological 4-manifold can be computed using the signature and the Rochlin invariant of the boundary with respect to the induced spin structure [FQ90, Proposition 10.2B]. To avoid the Rochlin invariant and having to specify spin structures, we instead glue the 0-trace of K , denoted by X_K , to W , along the common boundary $S_0^3(K)$; call the result Y . Since X_K is smooth, we know that $\text{ks}(Y) = \text{ks}(W)$. Using the Mayer–Vietoris sequence,

$$H_1(S_0^3(K); \mathbb{Z}) \xrightarrow{\cong} H_1(X_K; \mathbb{Z}) \oplus H_1(W; \mathbb{Z}) \twoheadrightarrow H_1(Y; \mathbb{Z}) \longrightarrow 0,$$

where we used that $H_1(X_K; \mathbb{Z}) = 0$, we see that $H_1(Y; \mathbb{Z}) = 0$. Using the Mayer–Vietoris sequence again, noting that $H_3(Y; \mathbb{Z}) \cong H^1(Y; \mathbb{Z}) = 0$, and that the inclusion induced map $H_2(S_0^3(K); \mathbb{Z}) \rightarrow H_2(X_K; \mathbb{Z})$ is an isomorphism while $H_2(S_0^3(K); \mathbb{Z}) \rightarrow H_2(W; \mathbb{Z})$ is trivial, we can further conclude that $H_2(Y; \mathbb{Z}) \cong H_2(W; \mathbb{Z})$ and indeed that the intersection forms on Y and on W coincide. So we have that Y is spin, since it has even intersection form and trivial first homology. By the definition of an n -solution, the signature $\sigma(Y) = \sigma(W) = 0$. For this we can either use that the intersection form of Y agrees with that of W , or use Novikov additivity. Putting everything together, we see that $\text{ks}(W) = \text{ks}(Y) = 0$ by [FQ90, Proposition 10.2B]. Therefore, the connected sum of W with sufficiently many copies of $S^2 \times S^2$ is smoothable [FQ90, Theorem 8.6] (see also [FNOP19b, Theorem 8.6; PR21, Section 21.4.5]). Choose $r \geq 0$ so that $W' = W \# rS^2 \times S^2$ is smoothable. Choose a smooth structure on W' . We will show that W' is a (smooth) n -solution.

As required, the manifold W' is smooth, compact, connected and oriented. We also know that inclusion induces an isomorphism $H_1(W; \mathbb{Z}) \cong H_1(W'; \mathbb{Z})$. So it remains only to check the conditions on the second homology. Let $\{L_i, D_i\}_{i=1}^k$ denote a basis for $H_2(W; \mathbb{Z})$ provided by the definition of an n -solution. By requiring that the connected sum operation is performed away from $\{L_i, D_i\}_i$, we may assume that each L_i and D_i also lies in W' . Let $\{A_i, B_i\}_{i=1}^r$ denote the standard basis for $H_2(rS^2 \times S^2; \mathbb{Z})$ given by the S^2 -factors, oriented so that each A_i intersects the corresponding B_i with positive sign. Of course each A_i and B_i has trivial normal bundle. We also know that $\pi_1(W') \cong \pi_1(W)$. Therefore, the set $\{L_i, D_i\}_{i=1}^k \cup \{A_i, B_i\}_{i=1}^r$ is a basis for $H_2(W'; \mathbb{Z})$ satisfying all the conditions of Definition 4.5, except that the elements of $\{L_i, D_i\}_i$ may not be smoothly embedded in W' .

In general, it may not be possible to change $\{L_i, D_i\}_i$ by a smooth ambient isotopy so that the result consists of smooth embeddings. However, by [FQ90, Theorem 8.1A] (see also [FQ90, p. 115]), we may modify each L_i and D_i by ambient isotopies and finger moves until the resulting immersions are smooth. The new intersections are paired by framed, embedded Whitney discs. Tube each L_i (or D_i) to itself, along one of each pair of Whitney arcs, as shown in Figure 4.1; call the resulting surfaces $\{L'_i, D'_i\}_i$. Then by construction each L'_i and D'_i is smoothly embedded in W' as needed. We also note that the intersections are as before, since the modification has occurred along arcs which can be assumed to have interiors disjoint from $\{L_i, D_i\}_i \cup \{A_i, B_i\}_i$ and one another. Of course, the genera of the surfaces have increased so we have to ensure that the new elements of the fundamental group still map




Figure 4.1: Left: The result of performing a finger move on L_i . The Whitney disc is shaded in grey. Right: The surface L'_i is produced after tubing along a Whitney arc. The blue curves show the new generators of $\pi_1(L'_i)$, denoted by μ and λ , compared to $\pi_1(L_i)$.

to $\pi_1(W)^{(n)}$ or $\pi_1(W)^{(n+1)}$, as appropriate. We consider the case of a surface L'_i – the case of a D'_i is similar. As shown in Figure 4.1, for each finger move, there are two new generators of the $\pi_1(L'_i)$ compared to $\pi_1(L_i)$, given by the curves μ and λ . However, both μ and λ are null-homotopic in W , the latter due to the existence of the Whitney disc and the former due to a meridional disc for L'_i . Therefore, the images of $\pi_1(L_i)$ and $\pi_1(L'_i)$ in $\pi_1(W)$ under the inclusion induced maps coincide. Now note that the set $\{L'_i, D'_i\}_{i=1}^k \cup \{A_i, B_i\}_{i=1}^r$ is a basis for $H_2(W'; \mathbb{Z})$ satisfying all the conditions of Definition 4.5. This uses the fact that each L'_i and D'_i is homologous in W' to the corresponding L_i and D_i , respectively, and completes the proof that W' is a smooth n -solution. \square

Consider the natural map $\Psi: \mathcal{F}_n \rightarrow \mathcal{F}_n^{\text{top}}$, where for each knot K the smooth concordance class of K is sent to the topological concordance class, i.e. induced by the map Ψ in (3.1). Then Proposition 4.7 has the following straightforward corollary.

Corollary 4.8. *For each $n \in \frac{1}{2}\mathbb{N}$, the map Ψ induces an isomorphism $\mathcal{F}_n/\mathcal{F}_{n+0.5} \cong \mathcal{F}_n^{\text{top}}/\mathcal{F}_{n+0.5}^{\text{top}}$.*

We now state the results mentioned in Section 4.1, as well as a few more open questions.

Theorem 4.9 ([COT03, COT04, CT07, CHL09, CHL11a]). *For each $n \in \mathbb{N}$, there is a subgroup of $\mathcal{F}_n/\mathcal{F}_{n.5}$ isomorphic to $\mathbb{Z}^\infty \oplus (\mathbb{Z}/2)^\infty$.*

For other related work, see also [Jia81, Liv99, DPR21]. By contrast, little is known about the “other half” of the filtration.

Question 4.10 (Open). *For some $n \in \mathbb{N}$, is the group $\mathcal{F}_{n.5}/\mathcal{F}_{n+1}$ nontrivial?*

For example, it was shown in [DMOP19] that genus one knots which are 0.5-solvable (equivalently, algebraically slice) are also 1-solvable. We note that there is an analogue of the solvable filtration for m -component (string) links, denoted by $\{\mathcal{F}_n^m\}_{n \in \frac{1}{2}\mathbb{N}}$, where for large enough m it is known that $\mathcal{F}_{n.5}^m/\mathcal{F}_{n+1}^m$ is nontrivial [Ott14].

Examples of knots lying in \mathcal{F}_n for large n are constructed using *satellite operations*. We will describe this further in Section 5. For now, we note that the examples from [COT03, COT04, CHL09, CHL11a] all have genus one. The examples from [DPR21] are cables of genus one knots, and have large Seifert genus and smooth slice genus, but may well have topological slice genus one. In general it is difficult to bound the topological slice genus of knots lying deep in the solvable filtration.

Question 4.11 (Open, Cha [Cha08, Remark 5.6]). *For arbitrary $n > 2$ and $g > 1$, does there exist a knot in \mathcal{F}_n with topological slice genus at least g ?*

The $n = 0$ and 1 cases can be shown using Levine–Tristram and Casson–Gordon signatures respectively, while the $n = 2$ case was shown by Cha–Miller–Powell in [CMP21].

Membership in the various levels of the solvable filtration can be obstructed using *von Neumann ρ -invariants*, defined by Cheeger and Gromov in [CG85]. For a detailed discussion, see [CHL11b, Section 5; CT07, Section 2; COT03, Section 2].

Theorem 4.12. *Let $K \subseteq S^3$ be a topologically slice knot. Then K lies in \mathcal{F}_n for every n . In other words, $\mathcal{T} \subseteq \bigcap_{n \in \frac{1}{2}\mathbb{N}} \mathcal{F}_n$.*

Proof. Since K is topologically slice, by Proposition 4.4 we can see that $K \in \mathcal{F}_n^{\text{top}}$ for every n by definition. Then by Proposition 4.7, we also know that $K \in \mathcal{F}_n$ for every n . \square

The above theorem implies that the solvable filtration is not useful for studying the structure of \mathcal{T} . However, there is another filtration which can be used for this purpose.

Definition 4.13 ([CHH13, Definition 5.1]). Let $K \subseteq S^3$ be a knot and $n \in \mathbb{N}$. We say that K is *n -positive* if $S_0^3(K) = \partial W$ where W is a smooth, compact, connected, oriented 4-manifold such that

- (i) the inclusion induces an isomorphism $\mathbb{Z} \cong H_1(S_0^3(K); \mathbb{Z}) \rightarrow H_1(W; \mathbb{Z})$;
- (ii) $H_2(W; \mathbb{Z})$ has a basis consisting of smoothly embedded, disjoint, closed, connected, oriented surfaces $\{S_i\}_{i=1}^k$, for some k , such that
 - (a) each S_i has normal bundle with Euler number $+1$; and
 - (b) for each i , we have $\pi_1(S_i) \subseteq \pi_1(W)^{(n)}$ with respect to the inclusion induced maps; and
- (iii) $\pi_1(W)$ is normally generated by the meridian $\mu_K \subseteq S_0^3(K)$.

The set of n -positive knots is denoted by \mathcal{P}_n and the manifold W is called an *n -positon*.

We say that K is *n -negative* if each surface S_i instead has normal bundle with Euler number -1 . The set of n -negative knots is denoted by \mathcal{N}_n and the manifold W is called an *n -negaton*.

Let $\mathcal{B}_n := \mathcal{P}_n \cap \mathcal{N}_n$. Knots in \mathcal{B}_n are said to be *n -bipolar*.

We leave it to the reader to show that, for each $n \in \mathbb{N}$, the sets \mathcal{P}_n and \mathcal{N}_n are submonoids of $\mathcal{C}^{\text{diff}}$, and \mathcal{B}_n is a subgroup of $\mathcal{C}^{\text{diff}}$ (Exercise Δ 13). As for the solvable filtration, note that condition (i) of Definition 4.13 and of Proposition 4.4 are the same. Indeed, now we further have the same condition (iii). Condition (ii) in Definition 4.13 is again a generalisation of condition (ii) in Proposition 4.4 – roughly speaking, if we assume that each surface S_i is a sphere, then one could perform a blow down operation on W along $\{S_i\}$, i.e. remove a neighbourhood of S_i diffeomorphic to the D^2 -bundle over S^2 with Euler number ± 1 , and then glue in B^4 , for each i , and the resulting 4-manifold with boundary $S_0^3(K)$ would have trivial second homology. The condition that $\pi_1(S_i) \subseteq \pi_1(W)^{(n)}$ as before measures how far away the homology class of S_i is from being represented by an immersed sphere: in the case that $\pi_1(S_i)$ is mapped to the trivial subgroup of $\pi_1(W)$, one could perform ambient surgery along half a symplectic basis of curves on S_i to transform it to an immersed sphere.

Definition 4.14. For each $n \in \mathbb{N}$ define $\mathcal{T}_n := \mathcal{T} \cap \mathcal{B}_n$. The corresponding filtration $\{\mathcal{T}_n\}_{n \in \mathbb{N}}$ is called the *bipolar filtration* of \mathcal{T} .

As desired, the bipolar filtration is highly nontrivial on \mathcal{T} , as seen in the theorem below.

Theorem 4.15 ([CK21], see also [CHH13, CH15]). *For each n , there is a subgroup of $\mathcal{T}_n/\mathcal{T}_{n+1}$ isomorphic to \mathbb{Z}^∞ .*

As before, the examples of knots lying in \mathcal{T}_n for large n used in the theorem above are constructed via a generalisation of the satellite operation, which we will describe in Section 5. As in the case of the solvable filtration, membership in the various levels of the positive, negative, and bipolar filtrations of $\mathcal{C}^{\text{diff}}$ can be obstructed using the von Neumann ρ -invariants. However, these are not sufficient to prove nontriviality of the bipolar filtration of \mathcal{T} . The proof of Theorem 4.15 combines von Neumann ρ -invariants with the Heegaard–Floer d -invariant. For more details, see [CHH13, CH15, CK21].

There are several open questions about the bipolar filtration, such as the following.

Question 4.16 (Open). Is there a (short) list of invariants characterising knots lying in \mathcal{B}_0 ? Note that there are explicit characterisations of knots in \mathcal{F}_0 and $\mathcal{F}_{0.5}$ (Exercise \circ 31).

Question 4.17 (Open). For $n \in \mathbb{N}$, is there a subgroup of $\mathcal{T}_n/\mathcal{T}_{n+1}$ isomorphic to $(\mathbb{Z}/2)^\infty$? (See Theorems 4.9 and 4.15.)

Question 4.18 (Open). For arbitrary $n \geq 0$ and $g > 1$, does there exist a knot in \mathcal{T}_n with smooth slice genus at least g ? (See Question 4.11.)

Unlike Question 4.11, the above appears to be open even in the case $n = 0$.

5. Satellite operations on knots

In this final section, we discuss satellite operations on knots. Recall that Whitehead doubling is a special case of the satellite operation, and satellite operations can be used to construct examples of knots deep in the solvable and bipolar filtrations.

Definition 5.1. A *pattern* P is a knot in the solid torus $S^1 \times D^2$. Given a knot K , called the *companion*, the *satellite knot* $P(K)$ is obtained by tying the solid torus into the knot K , in an appropriately untwisted manner, as shown in Figure 1.8. More precisely, we map $S^1 \times D^2$ to a tubular neighbourhood νK of K , so that $S^1 \times x$ for $x \in \partial D^2$ is mapped to the Seifert longitude of K . The image of P under this map is the satellite knot $P(K)$ by definition. Note that the knot $P(K)$ inherits an orientation from P and K .

The algebraic intersection number of P with a generic meridional disc of $S^1 \times D^2$ is called the *winding number* of P . For example, the winding number of the Whitehead doubling patterns Wh^\pm is zero.

The satellite operation is compatible with concordance, and yields well defined functions on the concordance groups, called *satellite operators* (see Exercise \triangle 15).

We next give an alternative view of the satellite construction. In this case, we begin with a knot $R \subseteq S^3$ and a curve $\eta \subseteq S^3 \setminus R$ so that η is unknotted when considered in $S^3 \supseteq S^3 \setminus R$. Then note that the complement in S^3 of a tubular neighbourhood $\nu\eta$ of η can be canonically identified with the solid torus, so $R \subseteq S^3 \setminus \nu\eta$ is a pattern. So, given any arbitrary knot K , we could construct the satellite knot with respect to this pattern. In this case, we denote the result by $R_\eta(K)$, and call it the *result of infection on R by K along η* . This procedure is depicted in Figure 5.1. There is a generalisation of the infection procedure where as input we can take not just knots but arbitrary *string links*. For more on this procedure, see [CO94, Section 1; Coc04, p. 385; CFT09, Section 2.2].

As previously indicated, the infection procedure can be used to construct knots in the various filtrations from the previous section. The key tool is the following proposition.

Proposition 5.2 ([COT04, Proposition 3.1; CHL09, Theorem 7.1; CHL11b, Proposition 2.7; CHH13, Proposition 3.3]). *Let $R \subseteq S^3$ be a knot and $\eta \subseteq S^3 \setminus R$ be a curve which is unknotted when considered in $S^3 \supseteq S^3 \setminus R$. Let $K \subseteq S^3$ be an arbitrary knot, and let $n, i \geq 0$ be arbitrary.*

1. *if $R, K \in \mathcal{F}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{F}_{n+i}$.*
2. *if $R, K \in \mathcal{P}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{P}_{n+i}$.*




Figure 5.1: On the left, we have a ribbon knot R with an unknot $\eta \subseteq S^3 \setminus R$. Given a knot K , we show on the right the result of infection on R by K along η .

3. if $R, K \in \mathcal{N}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{N}_{n+i}$.
4. if $R, K \in \mathcal{B}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{B}_{n+i}$.
5. if $R, K \in \mathcal{T}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{T}_{n+i}$.

We leave the proof for the ambitious reader (Exercise \square 27). Armed with this proposition, we are now able to construct knots deep in any of the solvable, positive, negative, or bipolar filtrations, by the following strategy. Let us consider the solvable filtration for concreteness and attempt to build a knot in \mathcal{F}_n for some given $n \in \mathbb{N}$. Start with any knot K with $\text{Arf}(K) = 0$, e.g. the connected sum of two copies of the right-handed trefoil knot. Recall that we know that $K \in \mathcal{F}_0$ by Exercise \circlearrowright 31. Then choose a ribbon knot R . By Theorem 4.12, we know that $R \in \bigcap_{i \in \frac{1}{2}\mathbb{N}} \mathcal{F}_i$. Find an unknotted $\eta \subseteq S^3 \setminus R$ such that $\text{lk}(R, \eta) = 0$. By definition of the linking number we know that $\eta \in \pi_1(S^3 \setminus R)^{(1)}$. So then $R_\eta(K) \in \mathcal{F}_1$ by Proposition 5.2. Then we iterate. In other words, to find a knot in \mathcal{F}_n , we need the n -fold iterated satellite $R_\eta(R_\eta(\dots(K)\dots))$. Note that these iterated satellites have bounded Seifert genus (Exercise \square 28) and we only needed the $i = 1$ case of the proposition.

A priori it might seem that satellites are quite special. However, by [CFT09, Proposition 1.7], any algebraically slice knot is concordant to a knot of the form $R_\eta(J)$, where J is a string link with pairwise linking numbers zero, the knot R is ribbon, and η is now an unlink, where each component has linking number zero with R . Therefore, the construction explained above is rather general.

We finish this section by surveying a few results and open questions about the effect of satellite operators on the knot concordance groups. We already saw questions about the Whitehead doubling satellite operator in Section 1. E.g. Conjecture 1.16 can be rephrased as asking whether the Whitehead doubling satellite operator is injective.

Question 5.3 (Open). Is there a pattern P with winding number $\neq \pm 1$ such that $P(K)$ is slice if and only if K is slice, in either category? More generally, such that $P(K)$ and $P(J)$ are concordant if and only if K and J are concordant?

Examples of winding number ± 1 satellite operators which are injective on \mathcal{C}^{top} were given in [CDR14]. From another perspective, one could ask when satellite operators are surjective on the concordance groups. For winding numbers $\neq \pm 1$ it is not difficult to find nonsurjective satellite operators using algebraic invariants [DR16, Proposition 3.1]. The first example of a nonsurjective satellite operator on $\mathcal{C}^{\text{diff}}$ with winding number ± 1 was given by A. Levine in [Lev16]. Other work has considered the invertibility of satellite operators, such as [DR16, MP18].

The existence of injective nonsurjective satellite operators is evidence towards the following conjecture. Further evidence was provided in [CHL11b] by showing that a large class of

winding number zero satellite operators, called *robust doubling operators*, are injective on large subgroups of $\mathcal{C}^{\text{diff}}$.

Conjecture 5.4 (Open, Cochran–Harvey–Leidy [CHL11b]). *The knot concordance groups have the structure of a fractal.*

We say a set has a fractal structure if it admits *self-similarities at arbitrarily small scales*, following [BGN03, Definition 3.1]. The results mentioned above show that infinite classes of winding number ± 1 satellite operators, as well as the robust doubling operators, are candidate self-similarities for \mathcal{C}^{top} and $\mathcal{C}^{\text{diff}}$. In order to understand the notion of “scale” one needs a suitable metric on the knot concordance groups. This has been studied in [CH18, CHP17, CHPR23]. One might also consider the behaviour of satellite operators under iteration, such as in [Ray15, Che20].

Since knots under concordance form a group and satellite operators are maps on the concordance groups, it is tempting to assume that they are homomorphisms. However, this is far from the case. In [Gom86], Gompf showed that the Whitehead doubling operator is not a homomorphism on $\mathcal{C}^{\text{diff}}$, despite being the zero map on \mathcal{C}^{top} . Modern tools from Heegaard–Floer homology can be used to obstruct satellite operators on $\mathcal{C}^{\text{diff}}$ from being homomorphisms [Lev16, Hed09]. In the topological category, obstructions can be obtained from Casson–Gordon invariants [Mil23]. Most generally, we have the following conjecture. (For a recent, partial resolution, see [JKM23].)

Conjecture 5.5 (Open, Hedden [Con16, Wor16]). *Let $P \subseteq S^1 \times D^2$ be a pattern. If the induced satellite operator $P: \mathcal{C}^{\text{diff}} \rightarrow \mathcal{C}^{\text{diff}}$ is a homomorphism, then it agrees with either the zero map, the identity map, or the reversal map (taking each knot to the concordance class of its reverse).*

Finally, there is substantial interest in understanding whether satellite operators preserve linear independence. We highlight two conjectures of Hedden and Pinzón-Caicedo.

Conjecture 5.6 (Open, Hedden–Pinzón-Caicedo [HPC21, Conjecture 2]). *The image of every nonconstant satellite operator on $\mathcal{C}^{\text{diff}}$ generates an infinite rank subgroup of $\mathcal{C}^{\text{diff}}$.*

The winding number $\neq 0$ case is not difficult to prove using algebraic invariants. [HPC21] provides a criterion guaranteeing that winding number zero satellite operators have infinite rank images, generalising [HK12, PC17].

Conjecture 5.7 (Open, Hedden–Pinzón-Caicedo [HPC21, Conjecture 3]). *For any nonconstant winding number zero operator P , there exists a knot K for which the set $\{P(nK) \mid n \in \mathbb{Z}\}$ has infinite rank in $\mathcal{C}^{\text{diff}}$.*

The above was partially confirmed in [DHMS22] using tools from Heegaard–Floer homology.

Epilogue

There are many aspects of the field of slice knots and knot concordance that we were unable to cover in these few lectures. So we end these notes with a brief, but surely incomplete, overview of those topics.

First, readers interested in more information on classical knot concordance should look at the excellent survey by Livingston [Liv05]. In particular, one will find there a much more detailed description of the algebraic concordance group and the Casson–Gordon invariants. Many recent developments in knot concordance, especially in the smooth setting, use Heegaard–Floer homology. Readers interested in those techniques should look at the excellent survey of Hom [Hom17]. Recent work has also leveraged *involutive* Heegaard–Floer homology [HM17]. Obstructions to sliceness can also be obtained from Khovanov homology [Ras10], and more generally \mathfrak{sl}_n -homologies [Lob09, LL16, LL19].

Quite a bit of recent research has considered knots in 3-manifolds other than S^3 and/or sliceness in 4-manifolds other than B^4 . A notable highlight is the proposed strategy of Freedman–Gompf–Morrison–Walker [FGMW10] to attack the 4-dimensional Poincaré conjecture: find a knot K and a homotopy 4-ball \mathcal{B} with $\partial\mathcal{B} = S^3$ such that K is smoothly slice in \mathcal{B} but not in B^4 . This would imply that \mathcal{B} is not diffeomorphic to B^4 , disproving the smooth 4-dimensional Poincaré conjecture. It is worth noting that the following weaker question is also open: does there exist a knot $K \subseteq S^3$ which is slice in an integer homology 4-ball but not in B^4 ? However there do exist *rationally slice* knots, such as the figure eight knot, i.e. knots that are slice in a rational homology ball, but not necessarily slice in B^4 . For more on this topic see e.g. [Kaw09, Cha07, HKPS22, HKP23].

More recent work of Manolescu and Piccirillo [MP21] explains how one can use sliceness of knots and links to address questions about exotic smooth structures on closed 4-manifolds other than S^4 . Other work concerning slicing knots in general 4-manifolds includes [Nor69, Yas91, Yas92, CT14, CN20, Rao20, MMP20, Pic20, KR21, KMRS21, MMRS22, MMSW23, HR23]. When studying concordance in more general 3-manifolds, one could also quotient out by the action of connected sum by knots in S^3 . This leads to the notion of *almost concordance*, previously called *piecewise linear I-equivalence* [Rol85; Hil12, Section 1.5; Cel18; FNOP19a; Yil18; NOPP19].

In light of Proposition 4.4 it is natural to ask to what extent the homeomorphism or homology cobordism class of the 0-surgery determines the concordance class of a knot. This is roughly the *Akbulut–Kirby* conjecture, For more on this, see e.g. [Kir97, Problem 1.19; KL99; CFHH13; Yas15; MP18].

We introduced ribbon discs in Section 1. There is a relative version called *ribbon concordance* in the smooth setting and *homotopy ribbon concordance* in the topological setting (see also [Exercise □ 21](#)). These are not necessarily symmetric relations. A flurry of recent work in the smooth setting [Zem19, LZ19, MZ21, DLVW22, Kan22] culminated in Agol showing that ribbon concordance is a partial order on knots [Ago22], confirming a conjecture of Gordon [Gor81]. It is still open whether homotopy ribbon concordance is a partial order as well [FP20, FKL⁺22].

We are also interested in investigating sliceness and concordance within standard families of knots. For example, we saw in Section 1 that the slice-ribbon conjecture holds for certain families. The families of algebraic knots [Rud76, Lit84, HKL12, CKP23], 2-bridge knots [CG86, Lis07, Mil18, FM16], pretzel knots [GJ11, Lec15, Lec12, Bry17, Lon14, Mil17a, Mil17b, KST20, KLS22], and (strongly) quasipositive knots [Rud93, Hay19, BF19] have received particular attention.

One may also study specific types of slice discs, e.g. those that have certain symmetries. There has been a renewed interest recently in *equivariant* sliceness and concordance [Nai97, CK99, DN06, BI22, DMS23, BC22, Mal22, MP23, DP23b, DPF23, DP23a].

Finally so far we have only discussed the existence of slice discs. One could equally well study the uniqueness question. In other words, can one quantify the number of slice discs for a given slice knot? Work in this area includes [Sch85, Akb91, CP21, SS22, JZ20, MP19, Akb22, HS21, LS22, Con22, DMS23].

Exercises

Introductory problems

Exercise Δ 1. Let $K \subseteq S^3$ be a knot, with tubular neighbourhood νK .

1. Show that $S^3 \setminus \nu K$ is a homology circle, i.e. $H_*(S^3 \setminus \nu K; \mathbb{Z}) \cong H_*(S^1; \mathbb{Z})$.
2. Show that $\pi_1(S^3 \setminus \nu K)$ is normally generated by an arbitrary meridian of K , i.e. it is generated by the set of conjugates of the meridian.

Do the above properties generalise to higher-dimensional knots $S^n \hookrightarrow S^{n+2}$? How about knots with arbitrary codimension?

Exercise Δ 2. Show that the connected sum operation is commutative and associative. In other words, given knots $J, K, L \subseteq S^3$, show that $J\#K$ is isotopic to $K\#J$ and $J\#(K\#L)$ is isotopic to $(J\#K)\#L$.

Exercise Δ 3. Prove Proposition 1.2: Let $K \subseteq S^3$ be a knot. Show that K is smoothly slice if and only if rK is smoothly slice if and only if \bar{K} is smoothly slice.

Note that the analogous statement also holds for topological sliceness.

Exercise Δ 4. Prove Proposition 1.3: If the knots $K, J \subseteq S^3$ are smoothly slice, then so is $K\#J$. As before, the analogous statement also holds for topological sliceness.




Figure 5.2: Some ribbon knots

Exercise Δ 5. Show that the knots in Figure 5.2 are ribbon.

Exercise Δ 6. Prove Proposition 1.6: For any knot $K \subseteq S^3$, the knot $K\#\bar{rK}$ is ribbon.

Exercise Δ 7. Prove Proposition 1.12: Let $K \subseteq S^3$ be a knot. The coned disc $\text{cone}(K) \subseteq \text{cone}(S^3) = B^4$ is locally flat if and only if K is the trivial knot.

Hint: Use the fact (without proof) from classical knot theory that $\pi_1(S^3 \setminus K) \cong \mathbb{Z}$ if and only if K is the trivial knot.

Exercise Δ 8. Let $K \subseteq S^3$ be a knot. Prove that if K is smoothly or topologically slice, then the same holds for the untwisted Whitehead doubles $\text{Wh}^\pm(K)$.

Exercise Δ 9. Prove Proposition 3.2: A knot $K \subseteq S^3$ is smoothly (resp. topologically) slice if and only if it is smoothly (resp. topologically) concordant to the unknot.

Exercise Δ 10. Prove Proposition 3.4: Smooth (resp. topological) concordance is an equivalence relation on knots.

Exercise Δ 11. Prove Proposition 3.6: In either $\mathcal{C}^{\text{diff}}$ or \mathcal{C}^{top} , the inverse of $[K]$ is the class of $-K := r\bar{K}$. The identity element in $\mathcal{C}^{\text{diff}}$ (resp. \mathcal{C}^{top}) is the class of the unknot, equivalently the class of any smoothly (resp. topologically) slice knot.

Exercise \triangle 12. Prove Proposition 3.7: A knot $K \subseteq S^3$ is said to be *negative amphichiral* if it is isotopic to \overline{rK} . Every negative amphichiral knot is order two in $\mathcal{C}^{\text{diff}}$ and \mathcal{C}^{top} . As a result, the figure eight knot is order two in $\mathcal{C}^{\text{diff}}$ and \mathcal{C}^{top} .

Exercise \triangle 13.

1. For each $n \in \frac{1}{2}\mathbb{N}$, show that the set \mathcal{F}_n is a subgroup of $\mathcal{C}^{\text{diff}}$.
2. For each $n \in \mathbb{N}$, show that the sets \mathcal{P}_n and \mathcal{N}_n are submonoids of $\mathcal{C}^{\text{diff}}$, and \mathcal{B}_n is a subgroup of $\mathcal{C}^{\text{diff}}$.

Exercise \triangle 14. Let $K \subseteq S^3$ be a smoothly slice knot. Show that K lies in \mathcal{T}_n for all n .

Exercise \triangle 15. Fix a pattern $P \subseteq S^1 \times D^2$. Show that the satellite operation is well defined on concordance on either category, i.e. there are well defined functions $P: \mathcal{C}^{\text{diff}} \rightarrow \mathcal{C}^{\text{diff}}$ and $P: \mathcal{C}^{\text{top}} \rightarrow \mathcal{C}^{\text{top}}$.

Exercise \triangle 16. Let $P \subseteq S^1 \times D^2$ be a pattern so that the induced satellite operator $P: \mathcal{C}^{\text{diff}} \rightarrow \mathcal{C}^{\text{diff}}$ (resp. $\mathcal{C}^{\text{top}} \rightarrow \mathcal{C}^{\text{top}}$) is a homomorphism. Show that $P(U)$ is smoothly (resp. topologically) slice, where U denotes the unknot.

Moderate problems

Exercise \square 17. Prove the hint from Exercise \triangle 7, that $\pi_1(S^3 \setminus K) \cong \mathbb{Z}$ if and only if K is the trivial knot.

Hint: Use Dehn's lemma.

Exercise \square 18. Learn enough classical knot theory to show that the right-handed trefoil, left-handed trefoil, and the figure eight from Figure 1.1 are nontrivial and distinct knots.

Hint: This will likely involve learning about some classical knot invariants, such as 3-colourability, the knot group, the Seifert genus, the signature, the Alexander polynomial, ...

Exercise \square 19. Fix $n \geq 4$. Prove that every smooth $S^1 \hookrightarrow S^n$ bounds a smoothly embedded disc in S^n .

Hint: The case $n = 4$ is the most challenging. Think about the types of singularities that arise in the generic case and try to get rid of them.

Exercise \square 20. Prove Proposition 1.5: A knot $K \subseteq S^3$ is ribbon if and only if it bounds a disc in S^3 with only ribbon singularities, i.e. singularities of the form shown in Figure 1.6.

Exercise \square 21. A knot $K \subseteq S^3$ is *homotopy ribbon* if there exists a topologically locally flat disc $D \subseteq B^4$ bounded by K such that the inclusion induced map $\pi_1(S^3 \setminus \nu K) \rightarrow \pi_1(B^4 \setminus \nu D)$ is surjective. Prove that ribbon implies homotopy ribbon.

Hint: Consider handle decompositions for ribbon disc complements.

Exercise \square 22. Prove Proposition 1.8: There exist smooth slice discs that are not ambiently isotopic (relative to the boundary) to any ribbon disc.

Hint: Begin with the standard smooth slice disc for the unknot. Use the fact that there exist 2-knots in S^4 with nonabelian fundamental group of the complement (the intrepid reader could try to prove the latter claim). Use Exercise \square 21. For a further challenge, construct examples of such discs bounded by nontrivial knots.

Exercise \square 23. Prove the trace embedding lemma (Lemma 2.4): Let $K \subseteq S^3$ be a knot. The 0-trace $X_0(K)$ admits a smooth (resp. locally collared) embedding into \mathbb{R}^4 if and only if K is smoothly (resp. topologically) slice.

Wonder whether a similar argument would apply to the n -traces of the knots, denoted by $X_n(K)$, obtained by attaching an n -framed 2-handle to B^3 along a knot $K \subseteq S^3 = \partial B^4$, for an arbitrary $n \in \mathbb{Z}$.

Hint: It is easier to prove the version of the trace embedding lemma in S^4 rather than \mathbb{R}^4 . Avoid using the smooth 4-dimensional Schoenflies conjecture (which is still open!) in the S^4 version by noting that the closure of the complement of a smoothly embedded 4-ball in S^4 is itself diffeomorphic to a 4-ball, by Palais's disc theorem.

Exercise □ 24. Prove Proposition 3.3: Two knots $K, J \subseteq S^3$ are smoothly (resp. topologically) concordant if and only if $K \# r\bar{J}$ is smoothly (resp. topologically) slice.

Hint: It is possible to isotope a concordance so that it contains a straight arc, i.e. one of the form $pt \times [0, 1] \subseteq S^3 \times [0, 1]$ (why?). Remove an open tubular neighbourhood of this arc. For the other direction, build a concordance between $K \# r\bar{J} \# J$ and J , and use Exercises 6 and 9.

Exercise □ 25. Show that isotopy classes of knots do **not** form a group under connected sum. They do however form a monoid.

Exercise □ 26. Reprove Proposition 4.4 using the trace embedding lemma (Lemma 2.4).

Exercise □ 27. Prove Proposition 5.2: Let $R \subseteq S^3$ be a knot and $\eta \subseteq S^3 \setminus R$ be a curve which is unknotted when considered in $S^3 \supseteq S^3 \setminus R$. Let $K \subseteq S^3$ be an arbitrary knot, and let $n, i \geq 0$ be arbitrary

1. if $R, K \in \mathcal{F}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{F}_{n+i}$.
2. if $R, K \in \mathcal{P}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{P}_{n+i}$.
3. if $R, K \in \mathcal{N}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{N}_{n+i}$.
4. if $R, K \in \mathcal{B}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{B}_{n+i}$.
5. if $R, K \in \mathcal{T}_n$ and $\eta \in \pi_1(S^3 \setminus R)^{(i)}$, then $R_\eta(K) \in \mathcal{T}_{n+i}$.

Exercise □ 28. Let $R \subseteq S^3$ be an arbitrary knot and $\eta \subseteq S^3 \setminus R$ be a curve which is unknotted when considered in $S^3 \supseteq S^3 \setminus R$. Assume that $lk(R, \eta) = 0$. Show that the Seifert genus of the knots produced by infection on R along η is bounded. In other words, there exists g , such that for any knot K , the Seifert genus of $R_\eta(K)$ is at most g . Come up with a candidate g .

Challenge problems

Exercise ○ 29. See Exercise □ 17. Are there any restrictions on the fundamental group of a slice/ribbon knot? In other words, is every knot group the knot group of a slice/ribbon knot?

Hint: Consider the Alexander module.

Exercise ○ 30. Prove that every smooth 2-knot is smoothly slice, i.e. every smooth $S^2 \hookrightarrow S^4$ bounds a smoothly embedded B^3 in B^5 . Prove that every topological 2-knot is topologically slice. This was originally proven by Kervaire [Ker65, Théorème III.6; Ker71, Theorem 1].

Hint: Use the fact that every 3-manifold is obtained by even-framed surgery on some link in S^3 .

Exercise ○ 31. Let $K \subseteq S^3$ be a knot.

1. Show that $K \in \mathcal{F}_0$ if and only if $\text{Arf}(K) = 0$.
2. Show that $K \in \mathcal{F}_{0.5}$ if and only if K is algebraically slice.
3. Show that if $K \in \mathcal{F}_{1.5}$ then all its Casson–Gordon sliceness obstructions vanish.

References

- [ACM⁺23] P. Aceto, N. A. Castro, M. Miller, J. Park, and A. Stipsicz, *Slice obstructions from genus bounds in definite 4-manifolds*, 2023. <https://arxiv.org/abs/2303.10587>.
- [Ago22] I. Agol, *Ribbon concordance of knots is a partial ordering*, *Commun. Am. Math. Soc.* **2** (2022), 374–379. MR4520779
- [Akb22] S. Akbulut, *Corks and exotic ribbons in B^4* , *Eur. J. Math.* **8** (2022), S494–S498. MR4507250
- [Akb91] S. Akbulut, *A fake compact contractible 4-manifold*, *J. Differ. Geom.* **33** (1991), no. 2, 335–356. MR1094459
- [AKPR21] P. Aceto, M. H. Kim, J. Park, and A. Ray, *Pretzel links, mutation, and the slice-ribbon conjecture*, *Math. Res. Lett.* **28** (2021), no. 4, 945–966. MR4344693
- [AT16] T. Abe and K. Tagami, *Fibered knots with the same 0-surgery and the slice-ribbon conjecture*, *Math. Res. Lett.* **23** (2016), no. 2, 303–323. MR3512887
- [BC22] K. Boyle and W. Chen, *Equivariant topological slice disks and negative amphichiral knots*, 2022. <https://arxiv.org/abs/2207.12593>.
- [BF19] M. Borodzik and P. Feller, *Up to topological concordance, links are strongly quasipositive*, *J. Math. Pures Appl.* **132** (2019), 273–279. MR4030255
- [BG96] Ž. Bižaca and R. E. Gompf, *Elliptic surfaces and some simple exotic \mathbf{R}^4 's*, *J. Differ. Geom.* **43** (1996), no. 3, 458–504. MR1412675
- [BGN03] L. Bartholdi, R. Grigorchuk, and V. Nekrashevych, *From fractal groups to fractal sets*, *Fractals in Graz 2001, 2003*, pp. 25–118. MR2091700
- [BI22] K. Boyle and A. Issa, *Equivariant 4-genera of strongly invertible and periodic knots*, *J. Topol.* **15** (2022), no. 3, 1635–1674. MR4461855
- [BKK⁺21] S. Behrens, B. Kalmár, M. H. Kim, M. Powell, and A. Ray (eds.), *The disc embedding theorem*, Oxford University Press, 2021. MR4519498
- [Bry17] K. Bryant, *Slice implies mutant ribbon for odd 5-stranded pretzel knots*, *Algebr. Geom. Topol.* **17** (2017), no. 6, 3621–3664. MR3709656
- [Cas86] A. J. Casson, *Three lectures on new-infinite constructions in 4-dimensional manifolds*, *À la recherche de la topologie perdue*, 1986, pp. 201–244. With an appendix by L. Siebenmann. MR900253
- [CDR14] T. D. Cochran, C. W. Davis, and A. Ray, *Injectivity of satellite operators in knot concordance*, *J. Topol.* **7** (2014), no. 4, 948–964. MR3286894
- [Cel18] D. Celoria, *On concordances in 3-manifolds*, *J. Topol.* **11** (2018), no. 1, 180–200. MR3784229
- [CFHH13] T. D. Cochran, B. D. Franklin, M. Hedden, and P. D. Horn, *Knot concordance and homology cobordism*, *Proc. Am. Math. Soc.* **141** (2013), no. 6, 2193–2208. MR3034445
- [CFT09] T. Cochran, S. Friedl, and P. Teichner, *New constructions of slice links*, *Comment. Math. Helv.* **84** (2009), no. 3, 617–638. MR2507256
- [CG78] A. J. Casson and C. McA. Gordon, *On slice knots in dimension three.*, *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976)*, Part 2, 1978, pp. 39–53. MR520521
- [CG85] J. Cheeger and M. Gromov, *Bounds on the von Neumann dimension of L^2 -cohomology and the Gauss-Bonnet theorem for open manifolds*, *J. Differ. Geom.* **21** (1985), no. 1, 1–34. MR806699
- [CG86] A. J. Casson and C. McA. Gordon, *Cobordism of classical knots*, *À la recherche de la topologie perdue*, 1986, pp. 181–199. With an appendix by P. M. Gilmer. MR900252
- [CG88] T. D. Cochran and R. E. Gompf, *Applications of Donaldson's theorems to classical knot concordance, homology 3-spheres and property P*, *Topology* **27** (1988), no. 4, 495–512. MR976591
- [CH15] T. D. Cochran and P. D. Horn, *Structure in the bipolar filtration of topologically slice knots*, *Algebr. Geom. Topol.* **15** (2015), no. 1, 415–428. MR3325742
- [CH18] T. Cochran and S. Harvey, *The geometry of the knot concordance space*, *Algebr. Geom. Topol.* **18** (2018), no. 5, 2509–2540. MR3848393
- [Cha07] J. C. Cha, *The structure of the rational concordance group of knots*, *Mem. Am. Math. Soc.* **189** (2007), no. 885, x+95. MR2343079
- [Cha08] J. C. Cha, *Topological minimal genus and L^2 -signatures*, *Algebr. Geom. Topol.* **8** (2008), no. 2, 885–909. MR2443100
- [Che20] W. Chen, *An infinite-rank summand from iterated Mazur pattern satellite knots*, 2020. <https://arxiv.org/abs/2010.11277>.
- [CHH13] T. D. Cochran, S. Harvey, and P. Horn, *Filtering smooth concordance classes of topologically slice knots*, *Geom. Topol.* **17** (2013), no. 4, 2103–2162. MR3109864
- [CHL09] T. D. Cochran, S. Harvey, and C. Leidy, *Knot concordance and higher-order Blanchfield duality*, *Geom. Topol.* **13** (2009), no. 3, 1419–1482. MR2496049
- [CHL11a] T. D. Cochran, S. Harvey, and C. Leidy, *2-torsion in the n -solvable filtration of the knot concordance group*, *Proc. Lond. Math. Soc.* **102** (2011), no. 2, 257–290. MR2769115
- [CHL11b] T. D. Cochran, S. Harvey, and C. Leidy, *Primary decomposition and the fractal nature of knot concordance*, *Math. Ann.* **351** (2011), no. 2, 443–508. MR2836668

- [CHP17] T. D. Cochran, S. Harvey, and M. Powell, *Grope metrics on the knot concordance set*, J. Topol. **10** (2017), no. 3, 669–699. MR3665407
- [CHPR23] T. D. Cochran, S. L. Harvey, M. Powell, and A. Ray, *Tower metrics on the smooth concordance set of knots*, 2023+. In preparation.
- [CK21] J. C. Cha and M. H. Kim, *The bipolar filtration of topologically slice knots*, Adv. Math. **388** (2021). MR4283760
- [CK99] J. C. Cha and K. H. Ko, *On equivariant slice knots*, Proc. Am. Math. Soc. **127** (1999), no. 7, 2175–2182. MR1605928
- [CKP23] A. Conway, M. H. Kim, and W. Politarczyk, *Nonslice linear combinations of iterated torus knots*, Algebr. Geom. Topol. **23** (2023), no. 2, 765–802. MR4587316
- [CL86] T. D. Cochran and W. B. R. Lickorish, *Unknotting information from 4-manifolds*, Trans. Am. Math. Soc. **297** (1986), no. 1, 125–142. MR849471
- [CMP21] J. C. Cha, A. N. Miller, and M. Powell, *Two-solvable and two-bipolar knots with large four-genera*, Math. Res. Lett. **28** (2021), no. 2, 331–382. MR4258974
- [CN20] A. Conway and M. Nagel, *Stably slice disks of links*, J. Topol. **13** (2020), no. 3, 1261–1301. MR4125756
- [CO94] T. D. Cochran and K. E. Orr, *Homology boundary links and Blanchfield forms: concordance classification and new tangle-theoretic constructions*, Topology **33** (1994), no. 3, 397–427. MR1286923
- [Coc04] T. D. Cochran, *Noncommutative knot theory*, Algebr. Geom. Topol. **4** (2004), 347–398. MR2077670
- [Con16] Conference organisers, *Problem List, Conference on 4-manifolds and knot concordance, Max Planck Institute for Mathematics*, 2016.
- [Con22] A. Conway, *Homotopy ribbon discs with a fixed group*, 2022. <https://arxiv.org/abs/2201.04465>.
- [COT03] T. D. Cochran, K. E. Orr, and P. Teichner, *Knot concordance, Whitney towers and L^2 -signatures*, Ann. Math. **157** (2003), no. 2, 433–519. MR1973052
- [COT04] T. D. Cochran, K. E. Orr, and P. Teichner, *Structure in the classical knot concordance group*, Comment. Math. Helv. **79** (2004), no. 1, 105–123. MR2031301
- [CP21] A. Conway and M. Powell, *Characterisation of homotopy ribbon discs*, Adv. Math. **391** (2021). MR4300918
- [CT07] T. D. Cochran and P. Teichner, *Knot concordance and von Neumann p -invariants*, Duke Math. J. **137** (2007), no. 2, 337–379. MR2309149
- [CT14] T. D. Cochran and E. Tweedy, *Positive links*, Algebr. Geom. Topol. **14** (2014), no. 4, 2259–2298. MR3331688
- [DHMS22] I. Dai, M. Hedden, A. Mallick, and M. Stoffregen, *Rank-expanding satellites, Whitehead doubles, and Heegaard Floer homology*, 2022. <https://arxiv.org/abs/2209.07512>.
- [DHST21] I. Dai, J. Hom, M. Stoffregen, and L. Truong, *More concordance homomorphisms from knot Floer homology*, Geom. Topol. **25** (2021), no. 1, 275–338. MR4226231
- [DKM+22] I. Dai, S. Kang, A. Mallick, J. Park, and M. Stoffregen, *The $(2, 1)$ -cable of the figure-eight knot is not smoothly slice*, 2022. <https://arxiv.org/abs/2207.14187>.
- [DLVVW22] A. Daemi, T. Lidman, D. S. Vela-Vick, and C.-M. M. Wong, *Ribbon homology cobordisms*, Adv. Math. **408** (2022). MR4467148
- [DMF92] S. De Michelis and M. H. Freedman, *Uncountably many exotic \mathbf{R}^4 's in standard 4-space*, J. Differ. Geom. **35** (1992), no. 1, 219–254. MR1152230
- [DMOP19] C. W. Davis, T. Martin, C. Otto, and J. Park, *Every genus one algebraically slice knot is 1-solvable*, Trans. Am. Math. Soc. **372** (2019), no. 5, 3063–3082. MR3988602
- [DMS23] I. Dai, A. Mallick, and M. Stoffregen, *Equivariant knots and knot Floer homology*, J. Topol. **16** (2023), no. 3, 1167–1236. MR4638003
- [DN06] J. F. Davis and S. Naik, *Alexander polynomials of equivariant slice and ribbon knots in S^3* , Trans. Am. Math. Soc. **358** (2006), no. 7, 2949–2964. MR2216254
- [Don83] S. K. Donaldson, *An application of gauge theory to four-dimensional topology*, J. Differ. Geom. **18** (1983), no. 2, 279–315. MR710056
- [Don86] S. K. Donaldson, *Connections, cohomology and the intersection forms of 4-manifolds*, J. Differ. Geom. **24** (1986), no. 3, 275–341. MR868974
- [Don87] S. K. Donaldson, *The orientation of Yang-Mills moduli spaces and 4-manifold topology*, J. Differ. Geom. **26** (1987), no. 3, 397–428. MR910015
- [DP23a] A. Di Prisa, *Equivariant algebraic concordance of strongly invertible knots*, 2023. <https://arxiv.org/abs/2303.11895>.
- [DP23b] A. Di Prisa, *The equivariant concordance group is not abelian*, Bull. Lond. Math. Soc. **55** (2023), no. 1, 502–507. MR4568356
- [DPF23] A. Di Prisa and G. Framba, *A new invariant of equivariant concordance and results on 2-bridge knots*, 2023. <https://arxiv.org/abs/2303.08794>.
- [DPR21] C. W. Davis, J. Park, and A. Ray, *Linear independence of cables in the knot concordance group*, Trans. Am. Math. Soc. **374** (2021), no. 6, 4449–4479. MR4251235
- [DR16] C. W. Davis and A. Ray, *Satellite operators as group actions on knot concordance*, Algebr. Geom. Topol. **16** (2016), no. 2, 945–969. MR3493412

- [End95] H. Endo, *Linear independence of topologically slice knots in the smooth cobordism group*, *Topology Appl.* **63** (1995), no. 3, 257–262. MR1334309
- [Fel16] P. Feller, *The degree of the Alexander polynomial is an upper bound for the topological slice genus*, *Geom. Topol.* **20** (2016), no. 3, 1763–1771. MR3523068
- [FGMW10] M. Freedman, R. Gompf, S. Morrison, and K. Walker, *Man and machine thinking about the smooth 4-dimensional Poincaré conjecture*, *Quantum Topol.* **1** (2010), no. 2, 171–208. MR2657647
- [FKL⁺22] S. Friedl, T. Kitayama, L. Lewark, M. Nagel, and M. Powell, *Homotopy ribbon concordance, Blanchfield pairings, and twisted Alexander polynomials*, *Can. J. Math.* **74** (2022), no. 4, 1137–1176. MR4464583
- [FM16] P. Feller and D. McCoy, *On 2-bridge knots with differing smooth and topological slice genera*, *Proc. Am. Math. Soc.* **144** (2016), no. 12, 5435–5442. MR3556284
- [FNOP19a] S. Friedl, M. Nagel, P. Orson, and M. Powell, *Satellites and concordance of knots in 3-manifolds*, *Trans. Am. Math. Soc.* **371** (2019), no. 4, 2279–2306. MR3896081
- [FNOP19b] S. Friedl, M. Nagel, P. Orson, and M. Powell, *A survey of the foundations of four-manifold theory in the topological category*, 2019. <https://arxiv.org/abs/1910.07372>.
- [Fox61] R. H. Fox, *Some problems in knot theory*, *Topology of 3-manifolds and related topics* (Proc. The Univ. of Georgia Institute, 1961), 1961, pp. 168–176. MR140100
- [Fox62] R. H. Fox, *A quick trip through knot theory*, *Topology of 3-manifolds and related topics* (Proc. The Univ. of Georgia Institute, 1961), 1962, pp. 120–167. MR0140099
- [FP20] S. Friedl and M. Powell, *Homotopy ribbon concordance and Alexander polynomials*, *Arch. Math.* **115** (2020), no. 6, 717–725. MR4179701
- [FQ90] M. H. Freedman and F. Quinn, *Topology of 4-manifolds*, *Princeton Mathematical Series*, vol. 39, Princeton University Press, 1990. MR1201584
- [Fre82] M. H. Freedman, *The topology of four-dimensional manifolds*, *J. Differ. Geom.* **17** (1982), no. 3, 357–453. MR679066
- [FS98] R. Fintushel and R. J. Stern, *Knots, links, and 4-manifolds*, *Invent. Math.* **134** (1998), no. 2, 363–400. MR1650308
- [FT86] M. H. Freedman and L. R. Taylor, *A universal smoothing of four-space*, *J. Differ. Geom.* **24** (1986), no. 1, 69–78. MR857376
- [GJ11] J. Greene and S. Jabuka, *The slice-ribbon conjecture for 3-stranded pretzel knots*, *Am. J. Math.* **133** (2011), no. 3, 555–580. MR2808326
- [Gom85] R. E. Gompf, *An infinite set of exotic \mathbf{R}^4 's*, *J. Differ. Geom.* **21** (1985), no. 2, 283–300. MR816673
- [Gom86] R. E. Gompf, *Smooth concordance of topologically slice knots*, *Topology* **25** (1986), no. 3, 353–373. MR842430
- [Gor81] C. McA. Gordon, *Ribbon concordance of knots in the 3-sphere*, *Math. Ann.* **257** (1981), no. 2, 157–170. MR634459
- [GS99] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, *Graduate Studies in Mathematics*, vol. 20, American Mathematical Society, 1999. MR1707327
- [GST10] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered knots and potential counterexamples to the property 2R and slice-ribbon conjectures*, *Geom. Topol.* **14** (2010), no. 4, 2305–2347. MR2740649
- [GT04] S. Garoufalidis and P. Teichner, *On knots with trivial Alexander polynomial*, *J. Differ. Geom.* **67** (2004), no. 1, 167–193. MR2153483
- [Hay19] K. Hayden, *Cross-sections of unknotted ribbon disks and algebraic curves*, *Compos. Math.* **155** (2019), no. 2, 413–423. MR3910456
- [Hed09] M. Hedden, *On knot Floer homology and cabling. II*, *Int. Math. Res. Not.* **12** (2009), 2248–2274. MR2511910
- [HHR16] M. A. Hill, M. J. Hopkins, and D. C. Ravenel, *On the nonexistence of elements of Kervaire invariant one*, *Ann. Math.* **184** (2016), no. 1, 1–262. MR3505179
- [Hil12] J. Hillman, *Algebraic invariants of links*, *Second, Series on Knots and Everything*, vol. 52, World Scientific, 2012. MR2931688
- [HK12] M. Hedden and P. Kirk, *Instantons, concordance, and Whitehead doubling*, *J. Differ. Geom.* **91** (2012), no. 2, 281–319. MR2971290
- [HKL12] M. Hedden, P. Kirk, and C. Livingston, *Non-slice linear combinations of algebraic knots*, *J. Eur. Math. Soc.* **14** (2012), no. 4, 1181–1208. MR2928848
- [HKL16] M. Hedden, S.-G. Kim, and C. Livingston, *Topologically slice knots of smooth concordance order two*, *J. Differ. Geom.* **102** (2016), no. 3, 353–393. MR3466802
- [HKP23] J. Hom, S. Kang, and J. Park, *Topologically and rationally slice knots*, 2023. <https://arxiv.org/abs/2304.06265>.
- [HKPS22] J. Hom, S. Kang, J. Park, and M. Stoffregen, *Linear independence of rationally slice knots*, *Geom. Topol.* **26** (2022), no. 7, 3143–3172. MR4540903
- [HLR12] M. Hedden, C. Livingston, and D. Ruberman, *Topologically slice knots with nontrivial Alexander polynomial*, *Adv. Math.* **231** (2012), no. 2, 913–939. MR2955197
- [HM17] K. Hendricks and C. Manolescu, *Involutive Heegaard Floer homology*, *Duke Math. J.* **166** (2017), no. 7, 1211–1299. MR3649355

- [Hom14] J. Hom, *The knot Floer complex and the smooth concordance group*, Comment. Math. Helv. **89** (2014), no. 3, 537–570. MR3260841
- [Hom15] J. Hom, *An infinite-rank summand of topologically slice knots*, Geom. Topol. **19** (2015), no. 2, 1063–1110. MR3336278
- [Hom17] J. Hom, *A survey on Heegaard Floer homology and concordance*, J. Knot Theory Ramifications **26** (2017), no. 2. MR3604497
- [Hom19] J. Hom, *Correction to the article An infinite-rank summand of topologically slice knots*, Geom. Topol. **23** (2019), no. 5, 2699–2700. MR4019902
- [HPC21] M. Hedden and J. Pinzón-Caicedo, *Satellites of infinite rank in the smooth concordance group*, Invent. Math. **225** (2021), no. 1, 131–157. MR4270665
- [HR23] M. Hedden and K. Raoux, *Knot Floer homology and relative adjunction inequalities*, Sel. Math., New Ser. **29** (2023), no. 1. MR4507976
- [HS21] K. Hayden and I. Sundberg, *Khovanov homology and exotic surfaces in the 4-ball*, 2021. <https://arxiv.org/abs/2108.04810>.
- [Jia81] B. J. Jiang, *A simple proof that the concordance group of algebraically slice knots is infinitely generated*, Proc. Am. Math. Soc. **83** (1981), no. 1, 189–192. MR620010
- [JKM23] R. Johanningsmeier, H. Kim, and A. N. Miller, *A partial resolution of Hedden’s conjecture on satellite homomorphisms*, 2023. <https://arxiv.org/abs/2308.06890>.
- [JZ20] A. Juhász and I. Zemke, *Distinguishing slice disks using knot Floer homology*, Sel. Math., New Ser. **26** (2020), no. 1. MR4045151
- [Kan22] S. Kang, *Link homology theories and ribbon concordances*, Quantum Topol. **13** (2022), no. 1, 183–205. MR4404800
- [Kaw09] A. Kawachi, *Rational-slice knots via strongly negative-amphicheiral knots*, Commun. Math. Res. **25** (2009), no. 2, 177–192. MR2554510
- [Kea73] C. Kearton, *Classification of simple knots by Blanchfield duality*, Bull. Am. Math. Soc. **79** (1973), 952–955. MR324706
- [Kea75a] C. Kearton, *Blanchfield duality and simple knots*, Trans. Am. Math. Soc. **202** (1975), 141–160. MR358796
- [Kea75b] C. Kearton, *Cobordism of knots and Blanchfield duality*, J. Lond. Math. Soc. **10** (1975), no. 4, 406–408. MR385873
- [Ker65] M. A. Kervaire, *Les nœuds de dimensions supérieures*, Bull. Soc. Math. Fr. **93** (1965), 225–271. MR189052
- [Ker71] M. A. Kervaire, *Knot cobordism in codimension two.*, Manifolds–Amsterdam 1970 (Proc. Nuffic Summer School), 1971, pp. 83–105. MR283786
- [Kir89] R. C. Kirby, *The topology of 4-manifolds*, Lecture Notes in Mathematics, vol. 1374, Springer, 1989. MR1001966
- [Kir97] R. Kirby, *Problems in low-dimensional topology*, Geometric topology (Athens, GA, 1993), 1997, pp. 35–473. MR1470751
- [KL22] T. Kim and C. Livingston, *Knot reversal acts non-trivially on the concordance group of topologically slice knots*, Sel. Math., New Ser. **28** (2022), no. 2. MR4363837
- [KL99] P. Kirk and C. Livingston, *Twisted knot polynomials: inversion, mutation and concordance*, Topology **38** (1999), no. 3, 663–671. MR1670424
- [KLS22] M. H. Kim, C. Lee, and M. Song, *Non-slice 3-stranded pretzel knots*, J. Knot Theory Ramifications **31** (2022), no. 3. MR4437814
- [KM63] M. A. Kervaire and J. W. Milnor, *Groups of homotopy spheres. I*, Ann. Math. **77** (1963), 504–537. MR148075
- [KMRS21] A. Kjuchukova, A. N. Miller, A. Ray, and S. Sakalli, *Slicing knots in definite 4-manifolds*, 2021. <https://arxiv.org/abs/2112.14596>.
- [KOPR21] M. H. Kim, P. Orson, J. Park, and A. Ray, *Open problems, The disc embedding theorem*, 2021, pp. 353–382. MR4519521
- [KR21] M. R. Klug and B. M. Ruppik, *Deep and shallow slice knots in 4-manifolds*, Proc. Am. Math. Soc., Ser. B **8** (2021), 204–218. MR4273166
- [KS77] R. C. Kirby and L. C. Siebenmann, *Foundational essays on topological manifolds, smoothings, and triangulations.*, Princeton University Press; University of Tokyo Press, 1977. With notes by John Milnor and Michael Atiyah. MR645390
- [KST20] K. Kishimoto, T. Shibuya, and T. Tsukamoto, *Sliceness of alternating pretzel knots and links*, Topology Appl. **282** (2020). MR4123275
- [Lec12] A. G. Lecuona, *On the slice-ribbon conjecture for Montesinos knots*, Trans. Am. Math. Soc. **364** (2012), no. 1, 233–285. MR2833583
- [Lec15] A. G. Lecuona, *On the slice-ribbon conjecture for pretzel knots*, Algebr. Geom. Topol. **15** (2015), no. 4, 2133–2173. MR3402337
- [Lev16] A. S. Levine, *Nonsurjective satellite operators and piecewise-linear concordance*, Forum Math. Sigma **4** (2016). MR3589337

- [Lev69a] J. Levine, *Invariants of knot cobordism*, Invent. Math. **8** (1969), 98–110. addendum, ibid. **8** (1969), **355**. MR253348
- [Lev69b] J. Levine, *Knot cobordism groups in codimension two*, Comment. Math. Helv. **44** (1969), 229–244. MR246314
- [Lis07] P. Lisca, *Lens spaces, rational balls and the ribbon conjecture*, Geom. Topol. **11** (2007), 429–472. MR2302495
- [Lit84] R. A. Litherland, *Cobordism of satellite knots*, Four-manifold theory (Durham, N.H., 1982), 1984, pp. 327–362. MR780587
- [Liv05] C. Livingston, *A survey of classical knot concordance*, Handbook of knot theory, 2005, pp. 319–347. MR2179265
- [Liv93] C. Livingston, *Knot theory*, Carus Mathematical Monographs, vol. 24, Mathematical Association of America, Washington, DC, 1993. MR1253070
- [Liv99] C. Livingston, *Order 2 algebraically slice knots*, Proceedings of the Kirbyfest (Berkeley, CA, 1998), 1999, pp. 335–342. MR1734416
- [LL16] L. Lewark and A. Lobb, *New quantum obstructions to sliceness*, Proc. Lond. Math. Soc. **112** (2016), no. 1, 81–114. MR3458146
- [LL19] L. Lewark and A. Lobb, *Upsilon-like concordance invariants from \mathfrak{sl}_n knot cohomology*, Geom. Topol. **23** (2019), no. 2, 745–780. MR3939052
- [Lob09] A. Lobb, *A slice genus lower bound from $\mathfrak{sl}(n)$ Khovanov-Rozansky homology*, Adv. Math. **222** (2009), no. 4, 1220–1276. MR2554935
- [Lon14] L. Long, *Slice Ribbon Conjecture, Pretzel Knots and Mutation*, Ph.D. Thesis, 2014.
- [LS22] R. Lipshitz and S. Sarkar, *Khovanov homology of strongly invertible knots and their quotients*, 2022. <https://arxiv.org/abs/2203.13895>.
- [LZ19] A. S. Levine and I. Zemke, *Khovanov homology and ribbon concordances*, Bull. Lond. Math. Soc. **51** (2019), no. 6, 1099–1103. MR4041014
- [Mal22] A. Mallick, *Knot Floer homology and surgery on equivariant knots*, 2022. <https://arxiv.org/abs/2201.07299>.
- [Mat02] Y. Matsumoto, *An introduction to Morse theory*, Translations of Mathematical Monographs, vol. 208, American Mathematical Society, 2002. Translated from the 1997 Japanese original by Kiki Hudson and Masahico Saito, Iwanami Series in Modern Mathematics. MR1873233
- [Mil17a] A. N. Miller, *Distinguishing mutant pretzel knots in concordance*, J. Knot Theory Ramifications **26** (2017), no. 7. MR3660096
- [Mil17b] A. N. Miller, *The topological sliceness of 3-strand pretzel knots*, Algebr. Geom. Topol. **17** (2017), no. 5, 3057–3079. MR3704252
- [Mil18] A. N. Miller, *A note on the topological sliceness of some 2-bridge knots*, Math. Proc. Camb. Philos. Soc. **164** (2018), no. 1, 185–191. MR3733246
- [Mil23] A. N. Miller, *Homomorphism obstructions for satellite maps*, Trans. Amer. Math. Soc., Ser. B **10** (2023), 220–240. MR4545198
- [Mil56] J. Milnor, *On manifolds homeomorphic to the 7-sphere*, Ann. Math. **64** (1956), 399–405. MR82103
- [Mil63] J. Milnor, *Morse theory.*, Princeton University Press, 1963. Based on lecture notes by M. Spivak and R. Wells. MR163331
- [Mil65] J. Milnor, *Lectures on the h-cobordism theorem.*, Princeton University Press, 1965. Notes by L. Siebenmann and J. Sondow. MR190942
- [Miy94] K. Miyazaki, *Nonsimple, ribbon fibered knots*, Trans. Am. Math. Soc. **341** (1994), no. 1, 1–44. MR1176509
- [MMP20] C. Manolescu, M. Marengon, and L. Piccirillo, *Relative genus bounds in indefinite four-manifolds*, 2020. <https://arxiv.org/abs/2012.12270>.
- [MMRS22] M. Marengon, A. N. Miller, A. Ray, and A. I. Stipsicz, *A note on surfaces in $\mathbb{C}P^2$ and $\mathbb{C}P^2 \# \mathbb{C}P^2$* , 2022. <https://arxiv.org/abs/2210.12486>.
- [MMSW23] C. Manolescu, M. Marengon, S. Sarkar, and M. Willis, *A generalization of Rasmussen’s invariant, with applications to surfaces in some four-manifolds*, Duke Math. J. **172** (2023), no. 2, 231–311. MR4541332
- [Moi77] E. E. Moise, *Geometric topology in dimensions 2 and 3*, Graduate Texts in Mathematics, Vol. 47, Springer, 1977. MR0488059
- [MP18] A. N. Miller and L. Piccirillo, *Knot traces and concordance*, J. Topol. **11** (2018), no. 1, 201–220. MR3784230
- [MP19] A. N. Miller and M. Powell, *Stabilization distance between surfaces*, Enseign. Math. **65** (2019), no. 3-4, 397–440. MR4113047
- [MP21] C. Manolescu and L. Piccirillo, *From zero surgeries to candidates for exotic definite four-manifolds*, 2021. <https://arxiv.org/abs/2102.04391>.
- [MP23] A. N. Miller and M. Powell, *Strongly invertible knots, equivariant slice genera, and an equivariant algebraic concordance group*, J. Lond. Math. Soc. **107** (2023), no. 6, 2025–2053. MR4598178
- [MZ21] M. Miller and I. Zemke, *Knot Floer homology and strongly homotopy-ribbon concordances*, Math. Res. Lett. **28** (2021), no. 3, 849–861. MR4270275

- [Nai97] S. Naik, *Equivariant concordance of knots in S^3* , KNOTS '96 (Tokyo), 1997, pp. 81–89. MR1664952
- [Nic11] L. Nicolaescu, *An invitation to Morse theory*, Second, Universitext, Springer, 2011. MR2883440
- [NOPP19] M. Nagel, P. Orson, J. Park, and M. Powell, *Smooth and topological almost concordance*, Int. Math. Res. Not. **23** (2019), 7324–7355. MR4039014
- [Nor69] R. A. Norman, *Dehn's lemma for certain 4-manifolds*, Invent. Math. **7** (1969), 143–147. MR246309
- [OSS17] P. S. Ozsváth, A. I. Stipsicz, and Z. Szabó, *Concordance homomorphisms from knot Floer homology*, Adv. Math. **315** (2017), 366–426. MR3667589
- [Ott14] C. Otto, *The (n) -solvable filtration of link concordance and Milnor's invariants*, Algebr. Geom. Topol. **14** (2014), no. 5, 2627–2654. MR3276843
- [PC17] J. Pinzón-Caicedo, *Independence of satellites of torus knots in the smooth concordance group*, Geom. Topol. **21** (2017), no. 6, 3191–3211. MR3692965
- [Pic20] J. Pichelmeyer, *Genera of knots in the complex projective plane*, J. Knot Theory Ramifications **29** (2020), no. 12. MR4193876
- [PR21] M. Powell and A. Ray, *The development of topological 4-manifold theory*, The disc embedding theorem, 2021, pp. 295–330. MR4519519
- [Qui82] F. Quinn, *Ends of maps. III. Dimensions 4 and 5*, J. Differ. Geom. **17** (1982), no. 3, 503–521. MR679069
- [Rao20] K. Raoux, *τ -invariants for knots in rational homology spheres*, Algebr. Geom. Topol. **20** (2020), no. 4, 1601–1640. MR4127080
- [Ras10] J. Rasmussen, *Khovanov homology and the slice genus*, Invent. Math. **182** (2010), no. 2, 419–447. MR2729272
- [Ray15] A. Ray, *Satellite operators with distinct iterates in smooth concordance*, Proc. Am. Math. Soc. **143** (2015), no. 11, 5005–5020. MR3391056
- [Rol85] D. Rolfsen, *Piecewise-linear 1-equivalence of links*, Low-dimensional topology (Chelwood Gate, 1982), 1985, pp. 161–178. MR827301
- [Rol90] D. Rolfsen, *Knots and links*, Mathematics Lecture Series, vol. 7, Publish or Perish, Inc., Houston, TX, 1990. Corrected reprint of the 1976 original. MR1277811
- [Rud76] L. Rudolph, *How independent are the knot-cobordism classes of links of plane curve singularities*, Notices Am. Math. Soc. **23** (1976), 410.
- [Rud93] L. Rudolph, *Quasipositivity as an obstruction to sliceness*, Bull. Am. Math. Soc. **29** (1993), no. 1, 51–59. MR1193540
- [Sch85] M. Scharlemann, *Smooth spheres in \mathbf{R}^4 with four critical points are standard*, Invent. Math. **79** (1985), no. 1, 125–141. MR774532
- [Sco05] A. Scorpan, *The wild world of 4-manifolds*, American Mathematical Society, 2005. MR2136212
- [SS22] I. Sundberg and J. Swann, *Relative Khovanov-Jacobsson classes*, Algebr. Geom. Topol. **22** (2022), no. 8, 3983–4008. MR4562563
- [Sta62] J. Stallings, *The piecewise-linear structure of Euclidean space*, Proc. Camb. Philos. Soc. **58** (1962), 481–488. MR149457
- [Sto77] N. W. Stoltzfus, *Unraveling the integral knot concordance group*, Mem. Am. Math. Soc. **12** (1977), no. 192, iv+91. MR467764
- [Tau87] C. H. Taubes, *Gauge theory on asymptotically periodic 4-manifolds*, J. Differ. Geom. **25** (1987), no. 3, 363–430. MR882829
- [Tro73] H. F. Trotter, *On S -equivalence of Seifert matrices*, Invent. Math. **20** (1973), 173–207. MR645546
- [Tro78] H. F. Trotter, *Knot module and Seifert matrices.*, Knot theory (Proc. Sem., Plans-sur-Bex, 1977), 1978, pp. 291–299. MR521739
- [Wor16] Workshop Organisers, *Final Report, Synchronizing Smooth and Topological 4-Manifolds*, 2016.
- [Yas15] K. Yasui, *Corks, exotic 4-manifolds and knot concordance*, 2015. <https://arxiv.org/abs/1505.02551>.
- [Yas91] A. Yasuhara, *$(2, 15)$ -torus knot is not slice in \mathbf{CP}^2* , Proc. Japan Acad., Ser. A **67** (1991), no. 10, 353–355. MR1151354
- [Yas92] A. Yasuhara, *On slice knots in the complex projective plane*, Rev. Mat. Univ. Complut. Madrid **5** (1992), no. 2-3, 255–276. MR1195083
- [Yil18] E. Z. Yildiz, *A note on knot concordance*, Algebr. Geom. Topol. **18** (2018), no. 5, 3119–3128. MR3848410
- [Zem19] I. Zemke, *Knot Floer homology obstructs ribbon concordance*, Ann. Math. **190** (2019), no. 3, 931–947. MR4024565